



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

The Country Heart Attack Prevention (CHAP) Project: implementation of an evidence based cardiac rehabilitation model of care for rural and remote patients: design paper

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-054558
Article Type:	Protocol
Date Submitted by the Author:	16-Jun-2021
Complete List of Authors:	<p>Beleigoli, Alline; Flinders University, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University Nicholls, Stephen ; Monash University Brown, Alex; SAHMRI, Indigenous Health Chew, Derek; Flinders Medical Centre, Department of Cardiovascular Medicine Beltrame, John; University of Adelaide, TQEH Discipline of Medicine Maeder, Anthony; Flinders University, College of Nursing & Health Sciences Maher, Carol; University of South Australia Versace, Vincent; Deakin University Geelong Australia, , Faculty of Health, School of Medicine Hendriks, Jeroen; Flinders University Tideman, Philip; Integrated Cardiovascular Clinical Network, Country Health SA Local Health Network, Flinders Medical Centre, Kaambwa, Billingsley; Flinders University Faculty of Medicine Nursing and Health Sciences, Health Economics Zeitz, Christopher; The University of Adelaide Faculty of Health and Medical Sciences Prichard, Ivanka; Flinders University Faculty of Social and Behavioural Sciences, School of Psychology Tavella, Rosanna; University of Adelaide, Medicine Tirimacco, Rosy; Integrated Cardiovascular Clinical Network SA Keech , Wendy; Health Translation SA Astley, C; Flinders University, Govin, Kay; Flinders University Faculty of Medicine Nursing and Health Sciences, College of Nursing and Health Sciences Nesbitt, Katie; Flinders University Du, Huiyun; Flinders University Champion, Stephanie; Flinders University Pinero de Plaza, Maria Alejandra; Flinders University College of Nursing and Health Sciences Lynch, Imelda; National Heart Foundation of Australia Poulsen, Vanessa; National Heart Foundation of Australia Ludlow, Marie; National Heart Foundation of Australia Wanguhu, Ken; Royal Australian College of General Practitioners Meyer, Hendrika; SA Health, Rural Support Services SA Health Krollig, Ali; Rural Support Services SA Gebremichael , Lemlem ; Flinders University College of Nursing and</p>

	Health Sciences Green , Chloe; Flinders University College of Nursing and Health Sciences Clark , Robyn ; Flinders University College of Nursing and Health Sciences
Keywords:	PREVENTIVE MEDICINE, CARDIOLOGY, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS

SCHOLARONE™
Manuscripts

TITLE PAGE

Title: The Country Heart Attack Prevention (CHAP) Project: implementation of an evidence-based cardiac rehabilitation model of care for rural and remote patients: design paper

Authors

1. Dr Alline Beleigoli, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
2. Professor Stephen J Nicholls, Monash Cardiovascular Research Centre, Victorian Heart Institute, Monash University
3. Professor Alex Brown, South Australian Health and Medical Research Institute Limited
4. Professor Derek P. Chew, College of Medicine and Public Health, Flinders University
5. Professor John Beltrame, The University of Adelaide, Central Adelaide Local Health Network
6. Professor Anthony Maeder, Flinders Digital Health Research Centre, Flinders University
7. Professor Carol Maher, University of South Australia
8. Associate Professor Vincent L. Versace, Deakin Rural Health, School of Medicine, Deakin University
9. Professor Jeroen M Hendriks, Caring Futures Institute, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University; Centre for Heart Rhythm Disorders, University of Adelaide and Royal Adelaide Hospital
10. Doctor Philip Tideman, Integrated Cardiovascular Clinical Network, Rural Support Service
11. Associate Professor Billingsley Kaambwa, College of Medicine and Public Health, Flinders
12. Associate Professor Christopher Zeitz, The Queen Elizabeth Hospital, the University of Adelaide
13. Associate Professor Ivanka Prichard, Caring Futures Institute, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
14. Assoc Prof Rosanna Tavella, The University of Adelaide, Central Adelaide Local Health Network
15. Rosy Tirimacco, Integrated Cardiovascular Clinical Network, Rural Support Service
16. Wendy Keech, Health Translation SA
17. Associate Professor Carolyn Astley, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
18. Kay Govin, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
19. Katie Nesbitt, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
20. Dr Huiyun Du, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
21. Dr Stephanie Champion, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

22. Dr Maria Alejandra Pinero de Plaza, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University; National Health and Medical Research Council (NHMRC) Centre of Research Excellence: Frailty and Healthy Ageing.

23. Imelda Lynch, National Heart Foundation of Australia

24. Vanessa Poulsen, National Heart Foundation of Australia

25. Dr Marie Ludlow, National Heart Foundation of Australia

26. Dr Ken Wanguhu, Royal Australian College of General Practitioners

27. Dr Hendrika Meyer, Rural Support Service SA Health

28. Ali Krollig, Rural Support Services SA Health

29. Dr Lemlem Gebremichael, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

30. Chloe Green, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

31. Prof Robyn Clark, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

Corresponding author:
Dr Alline Beleigoli, on behalf of the CHAP Project Team
Sturt North N214
GPO Box 2100, Adelaide 5001, South Australia
P:+61 8 82013303| alline.beleigoli@flinders.edu.au

ABSTRACT (300 WORDS)

INTRODUCTION

Despite extensive evidence of its benefits and recommendation by guidelines, cardiac rehabilitation (CR) remains highly underutilised with only 30-40% of patients attending a program after hospital discharge. We aim to implement and evaluate the Country Heart Attack Prevention (CHAP) model of care to improve CR attendance and completion for rural and remote participants.

METHODS AND ANALYSIS

CHAP will apply the Model for Large Scale Knowledge Translation to develop and implement a model of care to CR. Partnering with patients, clinicians and health service managers we will co-develop new approaches and refine/expand existing ones to address known barriers to CR attendance. A patient-centred approach will guide the co-design of telehealth options with patients and provide them with choices for CR modes of delivery. To increase referral rates, CHAP will promote endorsement of CR among clinicians and develop an electronic system that automatizes referrals of in-hospital eligible patients to CR. A business model that enables payment of CR delivered in primary care will enable sustainable access to CR. To promote CR quality improvement, professional development interventions and an accreditation program of CR services and programs will be developed. To evaluate 12-month CR attendance/completion (primary outcome), clinical and cost-effectiveness (secondary outcomes) between patients exposed and not exposed to CHAP, we will have a multi-design approach that encompasses a cohort study, a pre-post study and a comprehensive economic evaluation.

ETHICS AND DISSEMINATION

This study was approved by the Southern Adelaide Clinical Human Research Ethics Committee Research Committee (HREC/20/SAC/78), which approved a waiver of informed consent. Findings and dissemination to patients and clinicians will be through a public website, online educational sessions, and scientific publications. De-identified data will be available from the corresponding author on reasonable request.

Trial registration: Australia New Zealand Clinical Trials Registry (ANZCTR), ACTRN12621000222842, registered 03/03/2021.

Keywords: (3-10 words) *Cardiology, Preventive Medicine, Telemedicine*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- The study will apply the Model for Large Scale Knowledge Translation to partner with patients, health professionals and managers, co-develop and implement a model of care to increase attendance to cardiac rehabilitation in rural and remote areas of South Australia
- Applying a patient-centred care approach, the CHAP model of care will co-design web-based cardiac rehabilitation and allow patients to choose their preferred mode of delivery, evaluate patient-reported outcome and experiences measures (PROMs and PREMs)
- To evaluate the clinical effectiveness of the model of care, a multi-methods approach that includes a cohort study design, pre-post study design will be adopted
- A comprehensive economic analysis to evaluate the cost efficiencies to health services, and the cost-benefit to rural and remote patients and families will be performed.
- At this stage, the CHAP model of care may not suit the needs of Aboriginal and Torres Strait Islander peoples living in rural and remote areas of South Australia, and will warrant further cultural adaption.

INTRODUCTION

Cardiovascular diseases (CVD) are the number one cause of death globally, taking an estimated 17.9 million lives each year. CVD are a group of disorders of the heart and blood vessels and include coronary heart disease, cerebrovascular disease, rheumatic heart disease and other conditions such as heart failure and arrhythmias.¹ The rate of CVD secondary events globally is estimated to be between 26-50%.¹ CVD kills one Australian every 12 minutes and is a major cause of mortality accounting for 43,249 deaths in 2019.²

Of particular concern are those Australians living in rural and remote areas, who have 90% higher rates of CVD-related hospitalisation and 60% higher rates of CVD death than those in metropolitan areas.³ The further a person lives from a metropolitan centre, the greater their risk of hospitalisation and death from CVD.^{4 5 6} In South Australia, a southern, central state of Australia with 1.76 million inhabitants, approximately 23% of the population live in regional areas (i.e., towns, small cities and areas that lie beyond the capital city of Adelaide).⁷

Alarming, the prevalence of risk factors for heart disease in South Australia is higher than the national average⁸ and the age-standardised rate for hospital admissions related to heart disease was the third worst among Australian states in 2017 (120 per 10,000 persons) with some regional areas having mortality rates for coronary heart disease 18% higher than the state average.⁵

Cardiac Rehabilitation (CR) evidence translation gap

CR is the sum of interventions required to ensure physical, psychological and social recovery for patients after an acute cardiovascular event.⁹ Despite high levels of evidence supporting benefit and cost effectiveness of risk factor modification to reduce secondary CVD events through CR,¹⁰ Australian and international statistics, for the past 20 years report that only 20-50% of eligible patients attend.^{10 11 12}

1
2
3 26 Historically, CR has been delivered face-to-face to groups in acute hospitals or community
4
5 27 centres.¹³ Alternative methods for provision of CR are effective but few have been
6
7
8 28 implemented into practice.^{13 14} Therefore, we would argue that the evidence for strategies to
9
10 29 modify risk factors through CR is strong but the evidence for the mode of delivery has become
11
12 30 outdated.^{14 15} Research on the reasons for non-attendance have cited, cost and access issues ,
13
14 31 limited models of care to accommodate patients who wish to return to work and a lack of
15
16 32 person-centredness in service design and delivery.¹⁶ This particularly affects attendance of
17
18 33 women, people with disabilities, rural and remote dwellers, and culturally and linguistically
19
20 34 diverse populations.¹⁶
21
22
23
24
25

26 36 **Preliminary data and background of the Country Heart Attack Prevention (CHAP)**
27
28 37 **Project**
29

30 38 Preliminary data and background leading to the CHAP project have been developed over a
31
32 39 decade (2009-2018) through a longstanding partnership among local researchers, clinicians and
33
34 40 health service managers. Through this partnership, our team have summarised the evidence
35
36 41 around CR, identified local barriers, and implemented and measured performance of a series
37
38 42 of initiatives to improve CR referral, attendance and completion.^{4 9 12 14 16-20} Limited referrals
39
40 43 to CR, lack of patient-centred approaches and individualised choices, lack of sustainable
41
42 44 lifelong commitment to CR supported by primary care, and heterogeneous CR quality have
43
44 45 been identified as main barriers.^{16 20}
45
46
47
48
49

50
51 47 Limited referrals to CR
52

53 48 A range of factors influence clinicians' view about CR. Our preliminary work showed that
54
55 49 clinicians lacking cardiac qualifications may have limited knowledge and awareness of CR and
56
57 50 its benefits. Low agreement among clinicians on patients that are more likely to benefit from
58
59
60

CR, clinicians' personal lifestyle and health belief, and the knowledge on the availability and quality of local CR programs also impact on referral to CR.²⁰ The lack of standard administrative processes of referral and of reliable systems to support clinicians with these processes have also been identified as reasons for limited referrals to CR.^{16 20} To address this, the South Australian Department of Health's (SA Health) Integrated Cardiovascular Clinical Network (iCCnet SA), which is a state-wide provider of clinical network support and practice of evidence-based cardiac care and continuous quality improvement servicing regional, rural and remote areas of South Australia, implemented the Country Access to Cardiac Health (CATCH) program.^{17 18} This program includes a central referral process that receives CR referrals for patients discharged from public and private hospitals and directs them to a face-to-face or telephone-based CR service depending on patient's preferences and proximity to a CR community centre.¹⁸

Lack of patient-centred approaches and individualised choices

The CATCH program also implemented a telephone-based CR service that is offered to patients living at least 50km away from a CR service.¹⁸ The CATCH telephone program is aligned with findings from a systematic review conducted by members of our team that showed that individualised telehealth or home-based CR programs were effective alternative models that are comparable to the traditional hospital and community-based programs in terms of reducing CVD risk factors.^{14 18} These telehealth options are particularly important to Australians living in rural and remote areas where distances and lower availability of services amplify limitations to CR access.^{4 16 19 6} In an analysis conducted in 2015, regional and rural locations receiving the telephone-based CR program showed an increase in referral rates (97.3%), a reduction in all-cause readmissions (3.4% vs 1.8%, $p < 0.001$), and reduced length

1
2
3 75 of stay for those patients readmitted (3.5 vs 2.7 days, p=0.001) after the program
4
5 76 implementation.¹⁸
6
7
8 77

9
10 78 Lack of sustainable lifelong commitment to CR supported by primary care
11

12 79 Access to a cardiologist is a known enabler of CR uptake.²¹ To address the barrier of the
13
14 80 shortage of specialists in rural and remote areas throughout Australia,²² iCCnet has developed
15
16 81 a model of care (GP Hybrid network) that enables CR delivery through a strong collaboration
17
18 82 among general practitioners (GPs), practice nurses working in rural and remote areas in South
19
20 83 Australia and CR nurses working in the CATCH telephone-based service.²³ This service has
21
22 84 been evaluated and shown high attendance and completion rates, and enhanced clinical
23
24 85 outcomes for patients within 12 months, such as improved lipids and diabetes control. ²³ All
25
26 86 these measures of performance of the GP Hybrid network were enabled by the establishment
27
28 87 of the CATCH database by iCCnet. This CR dataset with standardised electronic data
29
30 88 collection serves to audit and feedback purposes to all public face-to-face services and the
31
32 89 telephone-based service in the state of South Australia.^{17 24}
33
34
35
36
37
38
39

40 91 Heterogeneous CR quality
41

42 92 Quality improvement of CR programs is associated with higher rates of attendance and
43
44 93 completion.²⁴ The Australian Cardiovascular Health and Rehabilitation Association (ACRA)
45
46 94 recommends five core components for quality delivery and outcomes of CR services - access
47
48 95 to services, assessment and monitoring, recovery and longer-term maintenance, lifestyle or
49
50 96 behavioural changes, medication adherence, evaluation and quality improvement.⁹ National
51
52 97 Quality indicators for CR were established by ACRA and the National Heart Foundation of
53
54 98 Australia in 2019.⁹ However, currently, there is no standardisation of the CR services and
55
56 99 programs with only a few meeting minimum duration standards.²⁵ Therefore, there is large
57
58
59
60

variation across services in terms of the content, measures, the number of sessions and length of the CR program.

Aims and Objectives

The CHAP project aims to develop and implement evidence-based approaches to address modifiable barriers to accessing and successfully completing CR in rural and remote areas of South Australia.

We hypothesise (H_1) that compared with patients not exposed to the CHAP model of care (i.e., receiving standard care) patients exposed to CHAP will have higher rates of attendance and completion of CR; higher rates of risk factor modification; higher rates of adherence to evidence-based pharmacotherapy; experience lower rates of morbidity and mortality at 30 days and 12 months and that the model will demonstrate cost effectiveness for both services and patients.

Specific objectives of the CHAP project are to:

- 1) Adopt a patient-centred approach by co-designing the CHAP model of care with patients family and clinicians and evaluate the impact on CR attendance and completion
- 2) Evaluate the clinical effectiveness of the CHAP model and
- 3) Evaluate the cost-effectiveness of the CHAP model

METHODS AND ANALYSIS

Study Design

The CHAP project will use a translation methodology (Model for Large Scale Knowledge Translation²⁶) to implement a model of care into practice. The model will be assessed through a multi-method approach which includes a pre-post and cohort study designs, and a comprehensive economic evaluation.

126

127 *The Model for Large Scale Knowledge Translation*

128 The Model for Large Scale Knowledge Translation²⁶ is a generalisable translation model that
129 intends to facilitate implementation of evidence into practice. It is particularly useful for large
130 scale collaborative projects involving researchers and clinical staff. The first three of the four
131 stages of the Model for Large Scale Knowledge Translation (1. Summarise the evidence; 2.
132 Identify local barriers, and 3. Measure performance) have been completed in the decade (2009-
133 2018) of preliminary work by researchers in our team and included 1) systematic review on the
134 effectiveness of alternative modes of delivery of CR;¹⁴ 2) identification of local barriers to CR
135 attendance;¹⁶ 3) the establishment of a central referral system (CATCH central referral system)
136 and telephone-based program (CATCH telephone program);^{17 18} and 4) clinical database
137 (CATCH database).¹⁷

138 The fourth stage of the model is the final and most complex one – *Ensure all patients receive*
139 *the intervention*. The model uses a “four Es” approach to improve reliability: engage, educate,
140 execute, and evaluate ²⁶ (Figure 1). This is the stage that will be developed in the CHAP project.
141 The CHAP model of care must fit each hospital or health networks current system, including
142 local culture and resources.

143 To complete this stage, the team will work closely with health networks and CR services
144 addressing barriers to implementing the CHAP model of care in rural and remote areas of the
145 state (Figure 2).

146 The evaluation of the CHAP model of care will comprise both clinical (Objective 2) and cost-
147 effectiveness (Objective 3) assessments, and will involve a multi-method approach:

- 148 I. A prospective cohort study comparing attendance and clinical outcomes of patients
149 exposed and non-exposed to CHAP - the cohort of patients eligible to CR will be
150 followed-up for 12 months from CHAP implementation and outcomes will be

compared between those exposed to the CHAP model and those not exposed to the model of care. The exposed and non-exposed groups will be age, sex and diagnostic-related groups (DRG) matched controls in a 1:3 ratio;²⁷

II. A pre-post study design comparing outcomes before (01/01/2016 to 30/04/2021) and after CHAP implementation (01/05/2021 to 30/04/2023). A comprehensive economic analysis will be performed to evaluate the cost efficiencies to health services, and the cost-benefit to rural and remote patients and families.

Setting

The CHAP model of care will be implemented within the six local health networks in regional, rural and remote South Australia through iCCnet.¹⁷ The iCCNet is part of the Rural Support Service (RSS) which is a business unit within the South Australian Health Service that provides flexible, responsive and innovative services to support the six regional local health networks. Regional health services are delivered to a population of 497, 003 people with the following age-group distribution: 0 to 4 years - 5.82%, 5 to 18 years -17.52%, 19 to 64 years - 56.68%, and 65 years and over - 19.98%. There are 23 CR services and the CATCH telephone service.²⁸

Participants

Eligible participants will be defined according to current international guideline criteria ²⁹ by DRGs based on the International Classifications of Diseases-10 (ICD-10) codes and the current guidelines for those who are eligible for referral to CR these include: acute myocardial infarction (I21, I21.0, I21.1, I21.2, I21.3, I21.4, I21.9), hypertensive heart disease (I11.0, I11.9, I13, I13.0, I13.2, I13.9), ischemic heart disease (I20-I25), arrhythmias (I48, I48.0, I48.1, I48.2, I48.3, I48.4, I48.9, I49.9), and/or heart failure (I50, I50.0, I50.1, I50.9).³⁰ Eligibility will also be defined based on DRG procedure codes for coronary artery surgery, percutaneous coronary intervention, and ventricular assist device.³¹

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

176 All adult (≥ 18 years) South Australians, including Aboriginal and Torres Strait Islander
177 peoples, discharged from hospital with a diagnosis that meets the eligibility criteria to CR and
178 living within the six local health networks in regional, rural and remote areas will be exposed
179 to the CHAP model of care. The comparison group (non-exposed to CHAP) will be comprised
180 by age, sex and DRG matched controls not living in these regional, rural and remote areas.

181 We acknowledge we are implementing standard guidelines that may not suit Aboriginal and
182 Torres Strait Islander peoples. A future project to culturally adapt the CHAP model of care
183 using appropriate Aboriginal and Torres Strait Islander translation methodology is warranted.

184
185 **Exposure- CHAP model of care**

186 To co-design and refine the CHAP model of care (Objective 1), we will address four different
187 barriers to CR attendance and completion previously identified (Figure 2)

188 *1) Limited number of referrals*

189 We will address the limited number of referrals by promoting CR endorsement among
190 clinicians and providing them with an electronic system that facilitates referrals.

191 To increase CR endorsement, we will co-design education resources with clinicians including
192 evidence-based pharmacotherapy resources, and a variety of paper-based, videos and web-
193 based education resources. This will ensure that health professionals' and CR patients' needs
194 are addressed regardless of technology literacy levels.³²

195 We will expand the capabilities of referral introduced by the CATCH central referral system
196 by developing an automatic electronic referral system that will streamline in-hospital referrals.
197 A central portal for patient discharge reports will provide contact information on patients that
198 have been discharged with applicable codes. Currently, these reports are run from several
199 different databases and have the potential to miss patients living in rural and remote areas. We

will first have automatic reports generated from information taken from the health department (SAH DataMart™). This will all run on the back end of the Integrated Point of Care Clinical System (iPOCCS™)³³ in which an excel document is generated daily, reporting on the previous 24h of discharged patients. A Uniform Resource Locator (URL) will be sent to designated users allowing them log on to iPOCCS and download the file.

2) *Lack of patient-centred approaches and individualised choices*

Face-to-face and telephone services are well established as modes of CR delivery in rural South Australian regions.¹⁸ Despite variability in duration and content, these CR programs are generally based on the National Heart Foundation of Australia²⁹ and ACRA guidelines⁹ being comprised of 10 modules (Initial Assessment and Goal Setting, Heart Education and Self-management, Exercise Training and Physical Activity, Healthy Eating and Weight management, Tobacco Cessation and Alcohol Reduction, Medication Education and Review, Managing Medical Risk Factors, Psychosocial Wellbeing, Activities of Daily Living, Reassessment and Completion) which are delivered over 6 to 12 weeks depending on patient's needs and decisions from services.

To expand choices of CR modes of delivery, CHAP will co-design with patients a web-based program that will be accessed through a personal log-in. The content of the portal will be based on the 10 National Heart Foundation of Australia modules.^{9 29} Individualised programs will be tailored according to patients' needs. The anticipated duration of each web module completion is 30 minutes per week. Those in the web-delivery mode will be required to read short texts, watch short videos (about 5 per module), set personal goals, self-report their dietary, physical activity, smoking and alcohol ingestion habits weekly through responding online questionnaires on the website. The web portal will have a function to collect physical activity data from wearables and smartphone apps patients may use. Patients will be able to interact with a CR nurse through a message system embedded in the web portal. They will be asked to

complete four Heart Health Risk Factor Assessments (baseline, 8-12 weeks after starting CR, 6 and 12-months after completing CR). These assessments will have an expected duration of 20 minutes and will be comprised of clinical and PROMS assessment using validated questionnaires that assess lifestyle habits, exercise capacity, quality of life, mental health (Table 1). Six and 12-month follow-up will be performed by the patient’s primary carer physician.

Table 1: Primary and Secondary outcomes, measurements and sources of data

Primary outcomes	Measures/ Instruments
Attendance/ Completion	Number of patients attended and completed CR among eligible ones/year
Secondary Outcomes	
CR Referral rates	Number of patients referred to CR/eligible to CR/year
30-day and 12-month hospital CVD-related hospital re-admission	NA
30-day and 12-month CVD-related mortality	NA
<i>Cardiovascular risk factor modification</i>	
Smoking	Self-reported number of cigarettes
Alcohol intake	Self-reported grams of alcohol consumed/week
Body mass index	Self-reported weight and height
Nutrition	CSIRO Healthy Diet Score ³⁴
Diabetes control	Hb A1C
Exercise, physical activity and functional capacity	6-minute walk test ³⁵ International Physical Activity Questionnaire-Short Form ³⁶ Specific Activity Questionnaire ³⁷
Medication adherence	Adherence in Chronic Diseases Scale ³⁸
<i>Patient-reported outcome measures (PROMs) and patient-reported experience measures (PREMs)</i>	
Anxiety	Cardiac anxiety questionnaire ³⁹
Depression	PHQ-2, PHQ-9 ⁴⁰
Quality of life	EQ-5D-5L ⁴¹

	Kansas City Cardiomyopathy Questionnaire-12 ⁴² (for patients with heart failure)
Health literacy	Brief Health Literacy Screening tool ⁴³
Patient satisfaction	Self-reported satisfaction

* NA Not Applicable; PHQ-2 and PHQ-9: the 2- and 9-question Patient Health Questionnaires, respectively; EQ-5D-5L: European Quality of Life- 5 Dimensions – 5 Levels

Patients and public involvement

The CHAP model of care is underpinned by a patient-centred approach. Patient-centred care or person-centred care is based on principles that those receiving healthcare must be treated with dignity and respect and be involved in all decisions about their health.⁴⁴ This approach is applied in this study through involvement of patients in the co-design of the CR web program and development of a model that allows patients to choose among different CR modes of delivery (face-to-face, telephone or web-based, or combinations of these modes- Figure 2).⁴⁴

⁴⁵ We will particularly promote the participation of older CR patients and people from cultural and linguistically diverse backgrounds to ensure that groups at increased risk of low technology and health literacy are involved in the co-designing of the web-based CR program. We anticipate this will increase patient adherence and engagement with the web-based CR program.⁴⁵

Updating the modalities of telehealth that are currently available will allow for greater flexibility of CR delivery which can improve access for patients in need. Importantly, this also provides an appropriate response to the COVID-19 pandemic when traditional CR service delivery is disrupted.^{46 47}

3) Lack of sustainable lifelong commitment to CR supported by primary care

CHAP project will build upon the GP Hybrid Model of CR delivery by creating a business model which allows primary care to support CR delivery and assessment of CR results (risk factor change) of patients undertaking telephone and web-based CR. The model will comprise

1
2
3 256 a ‘Heart Health Risk Factor’ pre-assessment at the beginning of the program (i.e. 1-2 weeks
4
5 257 post-discharge) which will last 75 minutes (60 minutes with a Practice Nurse and 15 minutes
6
7
8 258 with the GP). Re-assessments will occur at 8-12 weeks after the beginning of the program, and
9
10 259 6 months and 12 months post-completion of the CR program. These will comprise 45 to 60
11
12 260 minutes consultations with the practice nurse, and 15-minute consultations with the GP. This
13
14
15 261 has potential to enable long term support for heart health from GPs.

16
17 262 4) Heterogeneous CR quality

18
19 263 The CHAP project will seek adherence to the quality indicators through multiple online and
20
21 264 offline professional development programs targeting the practitioners, development of an
22
23 265 accreditation program and refinement of data collection tools on the national CR quality
24
25 266 indicators. In particular, the project will try to integrate objective patient-generated physical
26
27 267 activity tracking data to the CATCH CR database to enable objective monitoring of physical
28
29 268 activity during CR and personalised feedback.
30
31
32

33 269
34
35 270 **Outcomes**

36
37 271 *Primary outcome* – CR attendance rate defined as the number of patients commencing CR
38
39 272 among those eligible to CR.

40
41 273 *Secondary outcomes* - will compare risk factor modification, clinical outcomes, PROMS and
42
43 274 PREMS of the CHAP and non-CHAP groups. Measurements and data sources for the
44
45 275 secondary outcomes are presented in Table 1.
46
47
48

49 276
50
51 277 **Data Sources**

52
53 278 The CATCH CR database collects data uniformly across metropolitan, rural and remote areas
54
55 279 using a standardised electronic entry process, including electronic data capture, of patients
56
57 280 moving through public CR services from rural and remote to a metropolitan hospital or vice
58
59
60

281 versa.¹⁸ This database will be linked to the Integrated South Australian Integrated Activity
282 Collection (ISAAC), which is the SA Health administrative database, enabling follow-up and
283 assessment of hospital re-admissions and mortality.⁴⁸

284 **Bias**

285 To avoid information bias (i.e., systematic differences in the completeness or the accuracy of
286 data leading to differential misclassification of individuals regarding exposures or outcomes)
287 we are up-skilling clinicians to input data onto the CR databases as part of continuous quality
288 improvement processes. The goal is to fill the gap of scarcity of data on CR data in Australia
289 ⁴⁹ and achieve universal, uniform data entry in South Australian CR programs.

291 **Sample size**

292 The average state-wide attendance of CR from 2013-2015 was 28.3% (pre-CHAP estimate)¹⁰.
293 Based upon a 4% improvement in attendance (i.e., to 32.3%) as reported by the British
294 Association for Cardiovascular Prevention and Rehabilitation audit,⁵⁰ a total sample size of
295 1223 (CHAP) and 3669 (usual care, 1:3 matching) will produce a two-sided 95% confidence
296 interval for the difference in population proportions with a width that is equal to 0.06. Allowing
297 for a 20% dropout, we will need to observe 1529 patients (765 per year) in the group exposed
298 to CHAP that will be matched to 4587 patients in the group non-exposed to CHAP (usual care)
299 from the CATCH database. From 2015-2019, approximately 2000 patients were entered into
300 the CATCH CR database per year making this study highly feasible.

302 **Data analysis**

303 To evaluate the effectiveness of CHAP (Objective 2)

304 Unpaired *t*-tests, χ tests and Mann–Whitney *U* tests for non-normal data will be used for
305 comparisons for the pre-post analysis of the CHAP model components. Statistical significance

1
2
3 306 will be inferred if 2-tailed tests estimated $P<0.05$. Rates of referral, attendance, and completion
4
5 307 of CR per year; modification of risk factors; adherence to evidence-based pharmacotherapy
6
7 308 morbidity and mortality at 30 days and 12 months will be compared between CHAP and non-
8
9 309 CHAP attendees and presented as Odds Ratios and 95% Confidence Intervals. Matching will
10
11 310 be conducted by grouping all cases and controls into relevant groupings of Age, Sex and DRG
12
13 311 and analysed using the Mantel-Haenszel methods.⁵¹
14
15 312 Cox regression models adjusted for clinical and social factors (age, sex, the Index of Relative
16
17 313 Socio-Economic Advantage and Disadvantage (IRSAD) from the Australian Bureau of
18
19 314 Statistics' Socio-Economic Indexes for Areas-SEIFA score,⁵² accessibility and Remoteness
20
21 315 Index of Australia (ARIA+),⁵³ primary cardiac diagnosis, Charlson Index,⁵⁴ in-hospital length
22
23 316 of stay, prior heart failure, coronary disease, atrial fibrillation, revascularisation, malignancy)
24
25 317 will be performed to understand the association between exposure to the CHAP model and
26
27 318 attendance. For clinical outcomes (30-day and 12-month hospital CVD-related hospital re-
28
29 319 admission, 30-day and 12-month CVD-related mortality), the model will also be adjusted for
30
31 320 CR attendance.
32
33
34
35
36
37
38
39

40 322 *To evaluate the cost-effectiveness of CHAP (Objective 3)*
41
42

43 323 The economic evaluation of the CHAP model including the business model (explore the item
44
45 324 numbers that can be utilised to cover GP based CR and items that can be used to employ and
46
47 325 train practice nurses to assist with assessment, monitoring, collection of outcomes data and
48
49 326 triage to maximise GP consultation time) that involves primary care in CR delivery will
50
51 327 comprise: i) the cost efficiencies to health services, and ii) cost-benefit to rural and remote
52
53 328 patients and families.
54
55

56
57 329 *i. The cost efficiencies to the health services*
58
59
60

All CVD hospitalisations during the follow-up period will be included and assigned a cost, based on the diagnosis-related group category reference and length of stay. Costs of providing CR will be taken from a health system perspective and calculated over 1 year; this includes staffing, consumables, overheads, and capital expenses. Quality-of-life measures will be estimated at baseline, 6 and 12 months after CR completion, using the standard EQ-5D-5L.⁴¹ The utility is defined by the Department of Health and Ageing as a numerical value assigned by an individual to a preference for, or desirability of, a specific level of health status or a specific health outcome. Data on utilities and survival can be combined to calculate Quality-Adjusted Life-Year (QALY) by exposed and non-exposed groups using the Quality-Adjusted Survival Analysis (QASA) method.⁵⁵⁻⁵⁷ Incremental cost per QALY saved will be calculated as an incremental (rehabilitation v usual care) cost per patient divided by the incremental QALY per patient.⁵⁸

ii. Cost-benefit to rural and remote patients and families

Using an economic survey developed by the Western Australian Department of Health⁵⁹ we will quantify the economic impact of the CHAP model based on information at baseline including patient demographics such as household structure, employment status and the distances travelled to access services; the nature of the consultation, waiting time, whether the patient required time off work (and if so, how much), and the mode of transport used to travel to the hospital. The survey will include accompanying persons, and if the person required any time off work to do so. Finally, the survey will explore the patient's treatment options; for example, did they consider having an appointment in the city essential to their care? This data will be modelled to compare two scenarios for face-to-face CR; 1) if the patient travelled to the city to attend CR, or 2) if the patient travelled to the nearest rural CR program if this is within 60 minutes' drive (approximately 100 km). All results will be presented unadjusted for multiple comparisons and all analyses will be done on an

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

intention-to-treat basis. 95% confidence intervals for the difference in QALYs will be calculated with bootstrapping.⁵⁶

Time Frame of the Project

This will be a three-year project conducted over four calendar years July 1st, 2019-June 30th, 2022. Due to the impact of the COVID pandemic on the project implementation, the project will now be complete by 30th June 2023. Year 1 will include project staff employment and training, development of resources, education materials, study protocols, additional datasets, and Ethics approval. Recruitment and follow-up will be conducted over years 1-3, analysis and reporting in the last 6 months of Project Year 4.

ETHICS AND DISSEMINATION

Ethics approval and consent to participate

This study was approved by the Southern Adelaide Clinical Human Research Ethics Committee Research Committee on 28th, April 2020 (HREC/20/SAC/78). All consecutive patients who are eligible for CR according to the recommended DRG codes and referred for CR will have consent waiver to have their data used for this research.

Dissemination

Findings will be disseminated to patients and health professionals using various means and media. A public website (<https://chaproject.com.au>), online educational sessions with patients and health professionals working in CR, monthly newsletters sent to participants via email, social media (Twitter @CHAPproject, LinkedIn CHAP Project and Facebook), and scientific publications and conference presentations.

De-identified data will be available upon reasonable request.

Authors contributions: The study was conceived and designed by RC, SN, AB, DC, JB, AM, CM, VV, JH, PT, BK, CZ, IP, RT, RT, and WK. The first draft of the paper was written by AB with significant input from RC before receiving input from remaining authors; CA, KG, KN, HD, SC, MAPP, IL, VP, MR, KW, HM, AK, LG, and CG. All authors read and approved the final manuscript.

Funding: This study is supported by a peer-reviewed National Health and Medical Research Council (NHMRC Partnership Grant-GNT1169893) and by the following partners: Heart Foundation, Exercise and Sports Science Australia (ESSA), Australian Cardiac Rehabilitation Association (ACRA), Country SA Primary Health Network, Cardiac Society of Australia and New Zealand (CSANZ), Flinders Foundation, and Royal College of General Practitioners (RACGP), Astra Zeneca, Novartis, Health Translation SA, the Integrated Cardiac Clinical Network (iCCnet SA) and Country Health SA and the Enterprise Data and Information Department of Health and Wellbeing, SA Health.

Competing interests: The authors do not have competing interests.

Acknowledgements: VLV is funded by the Australian Government Department of Health through the Rural Health Multidisciplinary Training (RHMT) program.

JMH is supported by a Future Leader Fellowship provided by the National Heart Foundation of Australia.

CM is funded by a Medical Research Future Fund Investigator Grant (1193862).

1
2
3 405 **Legend of Figures and Tables**
4

5 406
6
7 407 **Figures**
8


9 408 Figure 1: Using the Model for Large Scale Knowledge Translation to develop CHAP

10
11 409 Figure 2: Addressing the barriers through the CHAP model of care (Exposure) according to the
12 410 4th Stage of the Model for Large Scale Knowledge Translation (‘Engage, Educate, Execute,
13 411 Evaluate’)
14 412

15
16
17 413

For peer review only


Figure 1: Using the Model for Large Scale Knowledge Translation to develop CHAP



(Pronovost PJ, Berenholtz SM, Needham DM. Translating evidence into practice: a model for large scale knowledge translation. *BMJ*. 2008;337:a1714.)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 2: Addressing the barriers through the CHAP model of care (Exposure) according to the 4th Stage of the Model for Large Scale Knowledge Translation (‘Engage, Educate, Execute, Evaluate’)



References

1. Cardiovascular diseases - WHO.: World Health Organization
2020 [Available from: https://www.who.int/health-topics/cardiovascular-diseases/#tab=tab_1
accessed 19/05/2021 2021.
2. Causes of Death 2019. ABS cat no 33030: Australian Bureau of Statistics (ABS), 2020.
3. Heart, stroke & vascular diseases: Australian Institute of Health & Welfare (AIHW), 2018.
4. Clark RA, Coffee N, Turner D, et al. Application of geographic modeling techniques to quantify spatial
access to health services before and after an acute cardiac event: the Cardiac Accessibility and
Remoteness Index for Australia (ARIA) project. *Circulation* 2012;125(16):2006-14. doi:
10.1161/circulationaha.111.083394 [published Online First: 2012/03/28]
5. Interactive Australian Heart Maps. Compare heart health indicators across Australia. : National
Heart Foundation of Australia.; 2021 [Available from:
[https://www.heartfoundation.org.au/health-professional-tools/interactive-heart-map-
australia](https://www.heartfoundation.org.au/health-professional-tools/interactive-heart-map-australia) accessed 16/06/2021 accessed 16/06/2021 2021.
6. Tideman P, Taylor AW, Janus E, et al. A comparison of Australian rural and metropolitan
cardiovascular risk and mortality: the Greater Green Triangle and North West Adelaide
population surveys. *BMJ Open* 2013;3(8):e003203. doi: 10.1136/bmjopen-2013-003203
7. Statistics ABo. 2016 Census Quick Stats. South Australia 2016 [Available from:
[https://quickstats.censusdata.abs.gov.au/census_services/getproduct/census/2016/quicksta
t/4?opendocument](https://quickstats.censusdata.abs.gov.au/census_services/getproduct/census/2016/quicksta
t/4?opendocument) accessed 16/06/2021 2021.
8. A time for action in South Australia. : National Heart Foundation of Australia
2018.
9. Woodruffe S, Neubeck L, Clark RA, et al. Australian Cardiovascular Health and Rehabilitation
Association (ACRA) core components of cardiovascular disease secondary prevention and
cardiac rehabilitation 2014. *Heart Lung Circ* 2015;24(5):430-41. doi:
10.1016/j.hlc.2014.12.008 [published Online First: 2015/02/01]
10. Anderson L, Thompson DR, Oldridge N, et al. Exercise-based cardiac rehabilitation for coronary
heart disease. *Cochrane Database Syst Rev* 2016 Jan 5;2016(1):CD001800 doi:
101002/14651858CD001800pub3 PMID: 26730878; PMCID: PMC6491180 2016
11. Redfern J, Hyun K, Chew DP, et al. Prescription of secondary prevention medications, lifestyle
advice, and referral to rehabilitation among acute coronary syndrome inpatients: results from
a large prospective audit in Australia and New Zealand. *Heart* 2014;100(16):1281-8. doi:
10.1136/heartjnl-2013-305296 [published Online First: 2014/06/09]
12. Astley CM, Chew DP, Keech W, et al. The Impact of Cardiac Rehabilitation and Secondary
Prevention Programs on 12-Month Clinical Outcomes: A Linked Data Analysis. *Heart Lung Circ*
2020;29(3):475-82. doi: 10.1016/j.hlc.2019.03.015 [published Online First: 2019/05/11]
13. Jelinek MV, Thompson DR, Ski C, et al. 40 years of cardiac rehabilitation and secondary prevention
in post-cardiac ischaemic patients. Are we still in the wilderness? *Int J Cardiol* 2015;179:153-
9. doi: 10.1016/j.ijcard.2014.10.154 [published Online First: 2014/12/03]
14. Clark RA, Conway A, Poulsen V, et al. Alternative models of cardiac rehabilitation: a systematic
review. *Eur J Prev Cardiol* 2015;22(1):35-74. doi: 10.1177/2047487313501093 [published
Online First: 2013/08/15]
15. West RR, Jones DA, Henderson AH. Rehabilitation after myocardial infarction trial (RAMIT): multi-
centre randomised controlled trial of comprehensive cardiac rehabilitation in patients
following acute myocardial infarction. *Heart* 2012;98(8):637-44. doi: 10.1136/heartjnl-2011-
300302 [published Online First: 2011/12/24]
16. Astley CM, Neubeck L, Gallagher R, et al. Cardiac Rehabilitation: Unraveling the Complexity of
Referral and Current Models of Delivery. *J Cardiovasc Nurs* 2017;32(3):236-43. doi:
10.1097/jcn.0000000000000332 [published Online First: 2016/04/15]

17. Tideman PA, Tirimacco R, Senior DP, et al. Impact of a regionalised clinical cardiac support network on mortality among rural patients with myocardial infarction. *Med J Aust* 2014;200(3):157-60. doi: 10.5694/mja13.10645 [published Online First: 2014/02/18]
18. Tideman P, Tirimacco R, Berry N, et al. Country access to cardiac health (CATCH) program: improving rehospitalisation rates and length of stay for cardiac rehabilitation using innovative telehealth delivery. *Heart, Lung and Circulation* 2015;24:S453. doi: 10.1016/j.hlc.2015.06.790
19. Clark RA, Wilkinson D, Coffee N, et al. Mapping services to support a patient's journey through evidence-based care pathways after a cardiac event. *Heart, Lung and Circulation* 2009;18:S266. doi: 10.1016/j.hlc.2009.05.656
20. Gallagher R, Neubeck L, Du H, et al. Facilitating or getting in the way? The effect of clinicians' knowledge, values and beliefs on referral and participation. *Eur J Prev Cardiol* 2016;23(11):1141-50. doi: 10.1177/2047487316630085 [published Online First: 2016/02/03]
21. Ghisi GL, Polyzotis P, Oh P, et al. Physician factors affecting cardiac rehabilitation referral and patient enrollment: a systematic review. *Clin Cardiol* 2013;36(6):323-35. doi: 10.1002/clc.22126 [published Online First: 2013/05/04]
22. Summit to address rural medical specialist shortage: Royal Australian College of General Practitioners 2018 [Available from: <https://www1.racgp.org.au/newsgp/professional/summit-to-address-rural-medical-specialist-shortage> accessed 18/01/2021 2021.
23. Clark C, Tirimacco R, Morfidis N, et al. Delivery of outpatient cardiac rehabilitation using a GP Hybrid/Telephone Program model. National Rural Health Alliance. Tasmania: Rural Health Org 2019.
24. Pack QR, Johnson LL, Barr LM, et al. Improving cardiac rehabilitation attendance and completion through quality improvement activities and a motivational program. *J Cardiopulm Rehabil Prev* 2013;33(3):153-9. doi: 10.1097/HCR.0b013e31828db386 [published Online First: 2013/04/19]
25. Gallagher R, Thomas E, Astley C, et al. Cardiac Rehabilitation Quality in Australia: Proposed National Indicators for Field-Testing. *Heart, Lung and Circulation* 2020;29(9):1273-77. doi: 10.1016/j.hlc.2020.02.014
26. Pronovost PJ, Berenholtz SM, Needham DM. Translating evidence into practice: a model for large scale knowledge translation. *BMJ* 2008;337:a1714. doi: 10.1136/bmj.a1714
27. Pearce N. Analysis of matched case-control studies. *BMJ* 2016;352:i969. doi: 10.1136/bmj.i969
28. Country SA Primary Health Network SA Health 2021 [Available from: <https://www.countrysaphn.com.au/> accessed 19/05/2021 2021.
29. Chew DP, Scott IA, Cullen L, et al. National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand: Australian clinical guidelines for the management of acute coronary syndromes 2016. *Med J Aust* 2016;205(3):128-33. doi: 10.5694/mja16.00368 [published Online First: 2016/07/29]
30. International Statistical Classification of Diseases and Related Health Problems 10th Revision: World Health Organization, 2016.
31. Independent Hospital Pricing Authority. International Classification of Diseases and Related Health Problems, 11th revision. Australian Modification (ICD-10-AM), 2019.
32. Norman CD, Skinner HA. eHealth Literacy: Essential Skills for Consumer Health in a Networked World. *J Med Internet Res* 2006;8(2):e9. doi: 10.2196/jmir.8.2.e9
33. Integrated Point of Care Clinical System Integrated Clinical Cardiovascular Network (iCCnet) SA 2021 [Available from: <https://www.iccnetsa.org.au/ipoccs.aspx> accessed 19/05/2021 2021.
34. Hendrie GA, Rebuli MA, Golley RK. Reliability and relative validity of a diet index score for adults derived from a self-reported short food survey. *Nutr Diet* 2017;74(3):291-97. doi: 10.1111/1747-0080.12303 [published Online First: 2017/07/22]
35. Bellet RN, Adams L, Morris NR. The 6-minute walk test in outpatient cardiac rehabilitation: validity, reliability and responsiveness—a systematic review. *Physiotherapy* 2012;98(4):277-86. doi: <https://doi.org/10.1016/j.physio.2011.11.003>

36. Craig CL, Marshall AL, Sjöström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95. doi: 10.1249/01.Mss.0000078924.61453.Fb [published Online First: 2003/08/06]
37. Rankin SL, Briffa TG, Morton AR, et al. A specific activity questionnaire to measure the functional capacity of cardiac patients. *Am J Cardiol* 1996;77(14):1220-3. doi: 10.1016/s0002-9149(97)89157-6 [published Online First: 1996/06/01]
38. Buszko K, Obońska K, Michalski P, et al. The Adherence Scale in Chronic Diseases (ASCD). The power of knowledge: the key to successful patient — health care provider cooperation. *Medical Research Journal* 2016;1(1):37-42. doi: 10.5603/MRJ.2016.0006
39. Eifert GH, Thompson RN, Zvolensky MJ, et al. The cardiac anxiety questionnaire: development and preliminary validity. *Behav Res Ther* 2000;38(10):1039-53. doi: 10.1016/s0005-7967(99)00132-1 [published Online First: 2000/09/27]
40. Arroll B, Goodyear-Smith F, Crengle S, et al. Validation of PHQ-2 and PHQ-9 to screen for major depression in the primary care population. *Ann Fam Med* 2010;8(4):348-53. doi: 10.1370/afm.1139 [published Online First: 2010/07/21]
41. McCaffrey N, Kaambwa B, Currow DC, et al. Health-related quality of life measured using the EQ-5D-5L: South Australian population norms. *Health Qual Life Outcomes* 2016;14(1):133. doi: 10.1186/s12955-016-0537-0 [published Online First: 2016/09/21]
42. Spertus JA, Jones PG. Development and Validation of a Short Version of the Kansas City Cardiomyopathy Questionnaire. *Circulation: Cardiovascular Quality and Outcomes* 2015;8(5):469-76. doi: 10.1161/CIRCOUTCOMES.115.001958
43. Haun J, Luther S, Dodd V, et al. Measurement variation across health literacy assessments: implications for assessment selection in research and practice. *J Health Commun* 2012;17 Suppl 3:141-59. doi: 10.1080/10810730.2012.712615 [published Online First: 2012/10/04]
44. Patient-Centred Care Improving Quality and Safety Through Partnerships With Patients and Consumers. *Australia Commission on Safety and Quality in Healthcare* 2011
45. Safety and Quality Improvement Guide Standard 2: Partnering with Consumers. In: Australia Co, ed. Sydney: Australia Commission on Safety and Quality in Healthcare, 2012.
46. Nicholls SJ, Nelson M, Astley C, et al. Optimising Secondary Prevention and Cardiac Rehabilitation for Atherosclerotic Cardiovascular Disease During the COVID-19 Pandemic: A Position Statement From the Cardiac Society of Australia and New Zealand (CSANZ)☆. *Heart, Lung and Circulation* 2020;29(7):e99-e104. doi: <https://doi.org/10.1016/j.hlc.2020.04.007>
47. Arnold RH, Tideman PA, Devlin GP, et al. Rural and Remote Cardiology During the COVID-19 Pandemic: Cardiac Society of Australia and New Zealand (CSANZ) Consensus Statement^{☆}. *Heart, Lung and Circulation* 2020;29(7):e88-e93. doi: 10.1016/j.hlc.2020.05.001
48. Admitted Patient Care Government of South Australia SA Health 2021 [accessed 18/01/2021 2021.
49. Driscoll A, Hinde S, Harrison A, et al. Estimating the health loss due to poor engagement with cardiac rehabilitation in Australia. *International Journal of Cardiology* 2020;317:7-12. doi: <https://doi.org/10.1016/j.ijcard.2020.04.088>
50. National Audit of Cardiac Rehabilitation (NACR) Quality and Outcomes Report 2018: British Heart Foundation, 2018.
51. Altman D. Practical Statistics for Medical Research. London: Chapman and Hall 1991.
52. Census of Population and Housing: Socio-economic Indexes for Areas (SEIFA): Australian Bureau of Statistics; 2016 [Available from: <https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/2033.0.55.001main+features12016> accessed 18/01/2021 2021.
53. Australian Statistical Geography Standard: Volume 5 - Remoteness Structure.: Australian Bureau of Statistics; 2018 [Available from: <https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.005> accessed 18/01/2021 2021.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

54. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40(5):373-83. doi: 10.1016/0021-9681(87)90171-8 [published Online First: 1987/01/01]

55. Glasziou PP, Simes RJ, Gelber RD. Quality adjusted survival analysis. *Stat Med* 1990;9(11):1259-76. doi: 10.1002/sim.4780091106 [published Online First: 1990/11/01]

56. Glasziou PP, Cole BF, Gelber RD, et al. Quality adjusted survival analysis with repeated quality of life measures. *Stat Med* 1998;17(11):1215-29. doi: 10.1002/(sici)1097-0258(19980615)17:11<1215::aid-sim844>3.0.co;2-y [published Online First: 1998/07/22]

57. Martin AJ, Glasziou PP, Simes RJ. A cardiovascular extension of the Health Measurement Questionnaire. *J Epidemiol Community Health* 1999;53(9):548-57. doi: 10.1136/jech.53.9.548 [published Online First: 1999/11/24]

58. Sullivan SD, Mauskopf JA, Augustovski F, et al. Budget impact analysis-principles of good practice: report of the ISPOR 2012 Budget Impact Analysis Good Practice II Task Force. *Value Health* 2014;17(1):5-14. doi: 10.1016/j.jval.2013.08.2291 [published Online First: 2014/01/21]

59. Rankin S, House A, Heath D. The real cost of accessing specialist medical services for rural and remote communities in Western Australia. . 5th National Rural Health Conference 1999.

COUNTRY HEART ATTACK PREVENTION (CHAP) PROJECT

**Stage 1: Summarise Evidence (2009-2018)**

Systematic review on effectiveness of alternative modes of CR delivery
 Establishment of the Integrated Cardiac Clinical Network (iCCnet)
 Implementation of the Country Access to Cardiac Health (CATCH) telephone-based CR service


**Stage 2: Identify Local Barriers (2009-2018)**

Main barriers identified:

- 1) Limited referrals to CR
- 2) Lack of patient-centred approaches and individualised choices
- 3) Lack of sustainable lifelong commitment to CR supported by primary care
- 4) Heterogeneous CR quality


**Stage 3: Measure Performance (2009-2018)**

Implementation of the Country Access to Cardiac Health (CATCH) database for audit and feedback
 Cardiac rehabilitation AUDIT 2013-2015 showing low referral, attendance and completion rates

Stage 4: Ensure all patients receive the CHAP model of care (2020-2023)

Using the Model for Large Scale Knowledge Translation to develop CHAP

169x325mm (66 x 66 DPI)



Addressing the barriers through the CHAP model of care (Exposure) according to the 4th Stage of the Model for Large Scale Knowledge Translation ('Engage, Educate, Execute, Evaluate')

402x442mm (38 x 38 DPI)

STROBE CHECKLIST

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	1 4
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	8-10
Objectives	3	State specific objectives, including any prespecified hypotheses	11
Methods			
Study design	4	Present key elements of study design early in the paper	12
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	13
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	14
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	14-18
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	18
Bias	9	Describe any efforts to address potential sources of bias	19
Study size	10	Explain how the study size was arrived at	19

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	20-21 Table 1
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	20-21
Results -Not applicable			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
<hr/>		
Discussion	Not applicable	
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
<hr/>		
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open

Implementation and prospective evaluation of the Country Heart Attack Prevention model of care to improve attendance and completion of cardiac rehabilitation for patients with cardiovascular diseases living in rural Australia: a study protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-054558.R1
Article Type:	Protocol
Date Submitted by the Author:	12-Jan-2022
Complete List of Authors:	<p>Beleigoli, Alline; Flinders University, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University Nicholls, Stephen ; Monash University Brown, Alex; SAHMRI, Indigenous Health Chew, Derek; Flinders Medical Centre, Department of Cardiovascular Medicine Beltrame, John; University of Adelaide, TQEH Discipline of Medicine Maeder, Anthony; Flinders University, College of Nursing & Health Sciences Maher, Carol; University of South Australia Versace, Vincent; Deakin University Geelong Australia, , Faculty of Health, School of Medicine Hendriks, Jeroen; Flinders University Tideman, Philip; Integrated Cardiovascular Clinical Network, Country Health SA Local Health Network, Flinders Medical Centre, Kaambwa, Billingsley; Flinders University Faculty of Medicine Nursing and Health Sciences, Health Economics Zeitz, Christopher; The University of Adelaide Faculty of Health and Medical Sciences Prichard, Ivanka; Flinders University Faculty of Social and Behavioural Sciences, School of Psychology Tavella, Rosanna; University of Adelaide, Medicine Tirimacco, Rosy; Integrated Cardiovascular Clinical Network SA Keech , Wendy; Health Translation SA Astley, C; Flinders University, Govin, Kay; Flinders University Faculty of Medicine Nursing and Health Sciences, College of Nursing and Health Sciences Nesbitt, Katie; Flinders University Du, Huiyun; Flinders University Champion, Stephanie; Flinders University Pinero de Plaza, Maria Alejandra; Flinders University College of Nursing and Health Sciences Lynch, Imelda; National Heart Foundation of Australia Poulsen, Vanessa; National Heart Foundation of Australia Ludlow, Marie; National Heart Foundation of Australia Wanguhu, Ken; Royal Australian College of General Practitioners Meyer, Hendrika; SA Health, Rural Support Services SA Health</p>

	Krollig, Ali; Rural Support Services SA Gebremichael , Lemlem ; Flinders University College of Nursing and Health Sciences Green , Chloe; Flinders University College of Nursing and Health Sciences Clark , Robyn ; Flinders University College of Nursing and Health Sciences
Primary Subject Heading:	Cardiovascular medicine
Secondary Subject Heading:	Health services research
Keywords:	PREVENTIVE MEDICINE, CARDIOLOGY, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS

SCHOLARONE™
Manuscripts

TITLE PAGE

Title: Implementation and prospective evaluation of the Country Heart Attack Prevention model of care to improve attendance and completion of cardiac rehabilitation for patients with cardiovascular diseases living in rural Australia: a study protocol

Authors

1. Dr Alline Beleigoli, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
2. Professor Stephen J Nicholls, Monash Cardiovascular Research Centre, Victorian Heart Institute, Monash University
3. Professor Alex Brown, South Australian Health and Medical Research Institute Limited
4. Professor Derek P. Chew, College of Medicine and Public Health, Flinders University
5. Professor John Beltrame, The University of Adelaide, Central Adelaide Local Health Network
6. Professor Anthony Maeder, Flinders Digital Health Research Centre, Flinders University
7. Professor Carol Maher, University of South Australia
8. Associate Professor Vincent L. Versace, Deakin Rural Health, School of Medicine, Deakin University
9. Professor Jeroen M Hendriks, Caring Futures Institute, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University; Centre for Heart Rhythm Disorders, University of Adelaide and Royal Adelaide Hospital
10. Doctor Philip Tideman, Integrated Cardiovascular Clinical Network, Rural Support Service
11. Associate Professor Billingsley Kaambwa, College of Medicine and Public Health, Flinders
12. Associate Professor Christopher Zeitz, The Queen Elizabeth Hospital, the University of Adelaide
13. Associate Professor Ivanka Prichard, Caring Futures Institute, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
14. Assoc Prof Rosanna Tavella, The University of Adelaide, Central Adelaide Local Health Network
15. Rosy Tirimacco, Integrated Cardiovascular Clinical Network, Rural Support Service
16. Wendy Keech, Health Translation SA
17. Associate Professor Carolyn Astley, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
18. Kay Govin, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
19. Katie Nesbitt, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
20. Dr Huiyun Du, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University
21. Dr Stephanie Champion, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

22. Dr Maria Alejandra Pinero de Plaza, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University; National Health and Medical Research Council (NHMRC) Centre of Research Excellence: Frailty and Healthy Ageing.

23. Imelda Lynch, National Heart Foundation of Australia

24. Vanessa Poulsen, National Heart Foundation of Australia

25. Dr Marie Ludlow, National Heart Foundation of Australia

26. Dr Ken Wanguhu, Royal Australian College of General Practitioners

27. Dr Hendrika Meyer, Rural Support Service SA Health

28. Ali Krollig, Rural Support Services SA Health

29. Dr Lemlem Gebremichael, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

30. Chloe Green, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

31. Prof Robyn Clark, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University

Corresponding author:
Dr Alline Beleigoli, on behalf of the CHAP Project Team
Sturt North N214
GPO Box 2100, Adelaide 5001, South Australia
P:+61 8 82013303| alline.beleigoli@flinders.edu.au

ABSTRACT (300 WORDS)

INTRODUCTION

Despite extensive evidence of its benefits and recommendation by guidelines, cardiac rehabilitation (CR) remains highly underutilised with only 30-40% of eligible patients participating. We aim to implement and evaluate the Country Heart Attack Prevention (CHAP) model of care to improve CR attendance and completion for rural and remote participants.

METHODS AND ANALYSIS

CHAP will apply the Model for Large Scale Knowledge Translation to develop and implement a model of care to CR in rural Australia. Partnering with patients, clinicians and health service managers, we will co-develop new approaches and refine/expand existing ones to address known barriers to CR attendance. CHAP will co-design a web-based CR program with patients expanding their choices to CR attendance. To increase referral rates, CHAP will promote endorsement of CR among clinicians and develop an electronic system that automatizes referrals of in-hospital eligible patients to CR. A business model that includes reimbursement of CR delivered in primary care by Medicare will enable sustainable access to CR. To promote CR quality improvement, professional development interventions and an accreditation program of CR services and programs will be developed. To evaluate 12-month CR attendance/completion (primary outcome), clinical and cost-effectiveness (secondary outcomes) between patients exposed (n=1223) and not exposed (n=3669) to CHAP, we will apply a multi-design approach that encompasses a prospective cohort study, a pre-post study and a comprehensive economic evaluation.

ETHICS AND DISSEMINATION

This study was approved by the Southern Adelaide Clinical Human Research Ethics Committee Research Committee (HREC/20/SAC/78), which approved a waiver of informed consent. Findings and dissemination to patients and clinicians will be through a public website, online educational sessions, and scientific publications. De-identified data will be available from the corresponding author on reasonable request.

Trial registration: Australia New Zealand Clinical Trials Registry (ANZCTR), ACTRN12621000222842, registered 03/03/2021.

Keywords: (3-10 words) *Cardiology, Preventive Medicine, Telemedicine*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- The study will apply the Model for Large Scale Knowledge Translation to implement evidence-based cardiac rehabilitation into practice.
- Co-design of telehealth-based cardiac rehabilitation with patients and evaluation of patient-reported outcomes and experiences demonstrate the patient-centred approach of CHAP.
- Data linkage between clinical and administrative databases will allow an accurate evaluation of the model’s clinical effectiveness.
- Analysis of the cost efficiencies to health services, and the cost-benefit to rural patients and families will be performed.
- The CHAP model of care is not culturally tailored to the needs of Aboriginal and Torres Strait Islander peoples living in rural South Australia.

INTRODUCTION

Cardiovascular diseases (CVD) are the number one cause of death globally, taking an estimated 17.9 million lives each year. CVD are a group of disorders of the heart and blood vessels and include coronary heart disease, cerebrovascular disease, rheumatic heart disease and other conditions such as heart failure and arrhythmias.¹ The rate of CVD secondary events globally is estimated to be between 26-50%.¹ CVD kills one Australian every 12 minutes and is a major cause of mortality accounting for 43,249 deaths in 2019.²

Of particular concern are those Australians living in rural and remote areas, who have 90% higher rates of CVD-related hospitalisation and 60% higher rates of CVD death than those in metropolitan areas.³ The further a person lives from a metropolitan centre, the greater their risk of hospitalisation and death from CVD.^{4 5 6} In South Australia, a southern, central state of Australia with 1.76 million inhabitants, approximately 23% of the population live in regional areas (i.e., towns, small cities and areas that lie beyond the capital city of Adelaide).⁷

Alarming, the prevalence of risk factors for heart disease in South Australia is higher than the national average⁸ and the age-standardised rate for hospital admissions related to heart disease was the third worst among Australian states in 2017 (120 per 10,000 persons) with some regional areas having mortality rates for coronary heart disease 18% higher than the state average.⁵

Cardiac Rehabilitation (CR) evidence translation gap

CR is the sum of interventions required to ensure physical, psychological and social recovery for patients after an acute cardiovascular event.⁹ Despite high levels of evidence supporting benefit and cost effectiveness of risk factor modification to reduce secondary CVD events through CR,¹⁰ Australian and international statistics, for the past 20 years report that only 20-50% of eligible patients participate.^{10 11 12}

1
2
3 26 Historically, CR has been delivered face-to-face to groups in acute hospitals or community
4
5 27 centres.¹³ Alternative methods for provision of CR are effective but few have been
6
7
8 28 implemented into practice.^{13 14 15} Therefore, we would argue that the evidence for strategies to
9
10 29 modify risk factors through CR is strong but the evidence for the mode of delivery has become
11
12 30 outdated.^{14 16} Research on the reasons for non-attendance have cited, cost and access issues ,
13
14 31 limited models of care to accommodate patients who wish to return to work and a lack of
15
16 32 person-centredness in service design and delivery.¹⁷ This particularly affects attendance of
17
18 33 women, people with disabilities, rural and remote dwellers, and culturally and linguistically
19
20 34 diverse populations.¹⁷ Moreover, the Covid-19 public health emergency has amplified the
21
22 35 barriers on accessing centre-based CR¹⁵.

23
24
25
26 36
27
28 37 **Preliminary data and background of the Country Heart Attack Prevention (CHAP)**
29
30 38 **Project**

31
32
33 39 Preliminary data and background leading to the CHAP project have been developed over a
34
35 40 decade (2009-2018) through a longstanding partnership among local researchers, clinicians and
36
37 41 health service managers. Through this partnership, our team have summarised the evidence
38
39 42 around CR, identified local barriers, and implemented and measured performance of a series
40
41 43 of initiatives to improve CR referral, attendance and completion.^{4 9 12 14 17-21} Limited referrals
42
43 44 to CR, lack of patient-centred approaches and individualised choices, lack of sustainable
44
45 45 lifelong commitment to CR supported by primary care, and heterogeneous CR quality have
46
47 46 been identified as main barriers.^{17 21}

48
49 47
50
51
52 48 Limited referrals to CR

53
54
55 49 A range of factors influence clinicians' view about CR. Our preliminary work showed that
56
57 50 clinicians lacking cardiac qualifications may have limited knowledge and awareness of CR and
58
59
60

its benefits.²¹ Low agreement among clinicians on patients that are more likely to benefit from CR, clinicians' personal lifestyle and health belief, and the knowledge on the availability and quality of local CR programs also impact on referral to CR.²¹ The lack of standard administrative processes of referral and of reliable systems to support clinicians with these processes have also been identified as reasons for limited referrals to CR.^{17 21} To address this, the South Australian Department of Health's (SA Health) Integrated Cardiovascular Clinical Network (iCCnet SA), which is a state-wide provider of clinical network support and practice of evidence-based cardiac care and continuous quality improvement servicing regional, rural and remote areas of South Australia, implemented the Country Access to Cardiac Health (CATCH) program.^{18 19} This program includes a central referral process that receives CR referrals for patients discharged from public and private hospitals and directs them to a face-to-face or telephone-based CR service depending on patient's preferences and proximity to a CR community centre.¹⁹ In a quasi-experimental pre-post analysis conducted in 2015, regional and rural locations receiving the telephone-based CR program showed an increase in referral rates (97.3%), a reduction in all-cause readmissions (3.4% vs 1.8%, $p<0.001$), and reduced length of stay for those patients readmitted (3.5 vs 2.7 days, $p=0.001$) after the program implementation.¹⁹

Lack of patient-centred approaches and individualised choices

The CATCH program also implemented a telephone-based CR service that is offered to patients living at least 50km away from a CR service.¹⁹ The CATCH telephone program is aligned with findings from a systematic review conducted by members of our team that showed that individualised telehealth or home-based CR programs were effective alternative models that are comparable to the traditional hospital and community-based programs in terms of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

reducing CVD risk factors.^{14 19} These telehealth options are particularly important to
Australians living in rural and remote areas where distances and lower availability of services
amplify limitations to CR access.^{4 17 20 6}

Lack of sustainable lifelong commitment to CR supported by primary care

Access to a cardiologist is a known enabler of CR uptake.²² To address the barrier of the
shortage of specialists in rural and remote areas throughout Australia,²³ iCCnet has developed
a model of care (GP Hybrid network) that enables CR delivery through a strong collaboration
among general practitioners (GPs), practice nurses working in rural and remote areas in South
Australia and CR nurses working in the CATCH telephone-based service.²⁴ This service has
been evaluated and shown high attendance and completion rates, and enhanced clinical
outcomes for patients within 12 months, such as improved lipids and diabetes control. ²⁴ All
these measures of performance of the GP Hybrid network were enabled by the establishment
of the CATCH database by iCCnet. This CR dataset with standardised electronic data
collection serves to audit and feedback purposes to all public face-to-face services and the
telephone-based service in the state of South Australia.^{18 25}

Heterogeneous CR quality

Quality improvement of CR programs is associated with higher rates of attendance and
completion.²⁵ The Australian Cardiovascular Health and Rehabilitation Association (ACRA)
recommends five core components for quality delivery and outcomes of CR services - access
to services, assessment and monitoring, recovery and longer-term maintenance, lifestyle or
behavioural changes, medication adherence, evaluation and quality improvement.⁹ National
Quality indicators for CR were established by ACRA and the National Heart Foundation of
Australia in 2019.⁹ However, currently, there is no standardisation of the CR services and

programs with only a few meeting minimum duration standards.²⁶ Therefore, there is large variation across services in terms of the content, measures, the number of sessions and length of the CR program.

Aims and Objectives

The CHAP project aims to develop and implement evidence-based approaches to address modifiable barriers to accessing and successfully completing CR in rural and remote areas of South Australia.

We hypothesise (H_1) that compared with patients not exposed to the CHAP model of care (i.e., receiving standard care) patients exposed to CHAP will have higher rates of attendance and completion of CR; higher rates of risk factor modification; higher rates of adherence to evidence-based pharmacotherapy; experience lower rates of morbidity and mortality at 30 days and 12 months and that the model will demonstrate cost effectiveness for both services and patients.

Specific objectives of the CHAP project are to:

- 1) Adopt a patient-centred approach by co-designing the CHAP model of care with patients family and clinicians and evaluate the impact on CR attendance and completion
- 2) Evaluate the clinical effectiveness of the CHAP model and
- 3) Evaluate the cost-effectiveness of the CHAP model

METHODS AND ANALYSIS

Research Design

The CHAP project is implementing a model of care within the real world. To understand and observe the various implementation activities and evaluate their different effects within this

126 complex context of non-linear and changing processes and elements, we will apply multiple
127 research methods.²⁷

128 The Model for Large Scale Knowledge Translation, which is particularly useful for large scale
129 collaborative projects involving researchers and clinical staff, is a generalisable translation
130 model that will be used to facilitate implementation of evidence-based interventions into
131 practice²⁸. The first three of the four stages of the Model for Large Scale Knowledge
132 Translation (1. Summarise the evidence; 2. Identify local barriers, and 3. Measure
133 performance) have been completed in the decade (2009-2018) of preliminary work by
134 researchers in our team and included 1) systematic review on the effectiveness of alternative
135 modes of delivery of CR;¹⁴ 2) identification of local barriers to CR attendance;¹⁷ 3) the
136 establishment of a central referral system (CATCH central referral system) and telephone-
137 based program (CATCH telephone program);^{18 19} and 4) clinical database (CATCH
138 database).¹⁸

139 The fourth stage of the model is the final and most complex one – *Ensure all patients receive*
140 *the intervention*. The model uses a “four Es” approach to improve reliability: engage, educate,
141 execute, and evaluate ²⁸ (Figure 1). This is the stage that will be developed in the CHAP project.
142 The CHAP model of care must consider the context, local culture and resources of each hospital
143 and health networks in the current system.

144 To complete this stage, the team will work closely with health networks and CR services
145 addressing barriers to implementing the CHAP model of care in rural and remote areas of the
146 state (Figure 2).

147 The evaluation of the CHAP model of care will comprise both clinical (Objective 2) and cost-
148 effectiveness (Objective 3) assessments, and will involve a multi-method approach:

149 I. A prospective cohort study comparing attendance and clinical outcomes of patients
150 exposed and non-exposed to CHAP - the cohort of patients eligible to CR will be

followed-up for 12 months from CHAP implementation and outcomes will be compared between those exposed to the CHAP model and those not exposed to the model of care. Attendance will be defined as attendance to ≥ 1 CR program session. Completion was defined as attendance to at least 75% of the CR sessions.²⁹ The exposed and non-exposed groups will be age, sex and diagnostic-related groups (DRG) matched controls in a 1:3 ratio;³⁰

II. A pre-post study design comparing outcomes before (01/01/2016 to 30/06/2021) and after CHAP implementation (01/07/2021 to 30/06/2023). A comprehensive economic analysis will be performed to evaluate the cost efficiencies to health services, and the cost-benefit to rural and remote patients and families.

Setting

The CHAP model of care will be implemented within the six local health networks in regional, rural and remote South Australia through iCCnet.¹⁸ The iCCNet is part of the Rural Support Service (RSS) which is a business unit within the South Australian Health Service that provides flexible, responsive and innovative services to support the six regional local health networks. Regional health services are delivered to a population of 497, 003 people with the following age-group distribution: 0 to 4 years - 5.82%, 5 to 18 years -17.52%, 19 to 64 years - 56.68%, and 65 years and over - 19.98%. There are 23 CR services and the CATCH telephone service.³¹

Participants

Eligible participants will be defined according to current international guideline criteria³² by DRGs based on the International Classifications of Diseases-10 (ICD-10) codes and the current guidelines for those who are eligible for referral to CR these include: acute myocardial infarction (I21, I21.0, I21.1, I21.2, I21.3, I21.4, I21.9), hypertensive heart disease (I11.0, I11.9, I13, I13.0, I13.2, I13.9), ischemic heart disease (I20-I25), arrhythmias (I48, I48.0, I48.1, I48.2,

1
2
3 176 I48.3, I48.4, I48.9, I49.9), and/or heart failure (I50, I50.0, I50.1, I50.9).³³ Eligibility will also
4
5 177 be defined based on DRG procedure codes for coronary artery surgery, percutaneous coronary
6
7 178 intervention, and ventricular assist device.³⁴
8
9
10 179 All adult (≥ 18 years) South Australians, including Aboriginal and Torres Strait Islander
11
12 180 peoples, discharged from hospital with a diagnosis that meets the eligibility criteria to CR
13
14 181 (acute myocardial infarction, hypertensive heart disease, ischemic heart disease, arrhythmias,
15
16 182 and/or heart failure, coronary artery surgery, percutaneous coronary intervention, and
17
18 183 ventricular assist device)³³ and living within the six local health networks in regional, rural
19
20 184 and remote areas will be exposed to the CHAP model of care. The comparison group (non-
21
22 185 exposed to CHAP) will be comprised by age, sex and DRG matched controls not living in these
23
24 186 regional, rural and remote areas.
25
26
27
28
29 187 With approximately 60% of the Aboriginal and Torres Strait Islander populations of South
30
31 188 Australia living in outside major cities in regional and remote areas,³⁵ it is likely that a number
32
33 189 of these individuals will be exposed to the CHAP model of care. However, we acknowledge
34
35 190 we are implementing standard guidelines that may not suit these groups. A future project to
36
37 191 culturally adapt the CHAP model of care using appropriate Aboriginal and Torres Strait
38
39 192 Islander translation methodology is warranted. There will be no stratification by
40
41 193 Aboriginal/indigenous status in this study.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

195 **Exposure- The CHAP model of care**

196 Exposure to the CHAP model of care will be defined as an individual, professional or service
197 receiving at least one of the approaches proposed to address four different local barriers to CR
198 attendance and completion that were previously identified by this team (Figure 2):

- 199 1) Limited number of referrals

200 We will address the limited number of referrals by promoting CR endorsement among
201 clinicians and providing them with an electronic system that facilitates referrals.

202 To increase CR endorsement, we will co-design education resources with clinicians including
203 evidence-based pharmacotherapy resources, and a variety of paper-based, videos and web-
204 based education resources. One example of this is the “Recommend video wall” which was
205 produced by the CHAP project through collection of testimonies from patients who undertook
206 CR and multidisciplinary teams working in hospitals and CR services who recommend CR³⁶.
207 This video wall has been displayed in events with professionals and patients and websites of
208 organisations such as ACRA and Heart Foundation as a tool to promote endorsement to CR.
209 This will ensure that health professionals’ and CR patients’ needs are addressed regardless of
210 technology literacy levels.³⁷

211 We will expand the capabilities of referral introduced by the CATCH central referral system
212 by developing an automatic electronic referral system that will streamline in-hospital referrals.
213 A central portal for patient discharge reports will provide contact information on patients that
214 have been discharged with applicable codes. Currently, these reports are run from several
215 different databases and have the potential to miss patients living in rural and remote areas. We
216 will first have automatic reports generated from information taken from the health department
217 (SAH DataMart™). This will all run on the back end of the Integrated Point of Care Clinical
218 System (iPOCCS™)³⁸ in which an excel document is generated daily, reporting on the previous
219 24h of discharged patients. A Uniform Resource Locator (URL) will be sent to designated users
220 allowing them log on to iPOCCS and download the file.

221 2) *Lack of patient-centred approaches and individualised choices*

222 Face-to-face and telephone services are well established as modes of CR delivery in rural
223 South Australian regions.¹⁹ Despite variability in duration and content, these CR programs are

1
2
3 224 generally based on the National Heart Foundation of Australia ³² and ACRA guidelines ⁹ being
4
5 225 comprised of 10 modules (Initial Assessment and Goal Setting, Heart Education and Self-
6
7 226 management, Exercise Training and Physical Activity, Healthy Eating and Weight
8
9 227 management, Tobacco Cessation and Alcohol Reduction, Medication Education and Review,
10
11 228 Managing Medical Risk Factors, Psychosocial Wellbeing, Activities of Daily Living,
12
13 229 Reassessment and Completion) which are delivered over 6 to 12 weeks depending on patient's
14
15 230 needs and decisions from services.
16
17 231 To expand choices of CR modes of delivery, CHAP will co-design with patients living in rural
18
19 232 areas a web-based program accessed through a personal log-in. The *User Experience* (UX)
20
21 233 *Design*,³⁹ an established co-design framework, will guide the engagement with end-users
22
23 234 through workshops and the co-design process from concept design to completion. A variety of
24
25 235 UX Design methods will be used interchangeably to plan, execute and evaluate the workshops.
26
27 236 These include cultural probes, heuristic evaluation, brainstorming, focus groups and concept
28
29 237 testing.^{39 40} The content of the portal will be based on the 10 National Heart Foundation of
30
31 238 Australia modules.^{9 32} Individualised programs will be tailored according to patients' needs.
32
33 239 The anticipated duration of each web module completion is 30 minutes per week.⁴⁰ Those in
34
35 240 the web-delivery mode will be required to read short texts, watch short videos (about 5 per
36
37 241 module), set personal goals, self-report their dietary, physical activity, smoking and alcohol
38
39 242 ingestion habits weekly through responding online questionnaires on the website. The web
40
41 243 portal will have a function to collect physical activity data from wearables and smartphone
42
43 244 apps patients may use. Patients will be able to interact with a CR nurse through a message
44
45 245 system embedded in the web portal. They will be asked to complete four Heart Health Risk
46
47 246 Factor Assessments (baseline, 8-12 weeks after starting CR, 6 and 12-months after completing
48
49 247 CR). These assessments will comprise clinical and PROMS assessment using validated
50
51 248 questionnaires that assess lifestyle habits, exercise capacity, quality of life, mental health
52
53
54
55
56
57
58
59
60

(Table 1). Six and 12-month follow-up will be performed by the patient's primary carer physician.

Table 1: Primary and Secondary outcomes, measurements and sources of data

Primary outcomes	Measures/ Instruments
Attendance/ Completion	Number of patients attending ≥ 1 session or completing 75% of all sessions among the ones eligible ones/year
Secondary Outcomes	
CR Referral rates	Number of patients referred to CR/eligible to CR/year
30-day and 12-month hospital CVD-related hospital re-admission	Assessed through the linkage between the South Australian CR database and SA Health administrative databases. Each cardiac admission will be counted as a single separation, with admissions involving transfer(s) merged as one. Readmission within 24 hours will not be counted as a new event.
30-day and 12-month CVD-related mortality	Assessed through the linkage between the South Australian CR database and SA Health administrative databases. CVD
30-day and 12-month Emergency Department (ED) presentation	Assessed through the linkage between the South Australian CR database and SA Health administrative databases. Each ED presentation resulting in a separation will be considered a single hospitalisation. New ED presentation within 24 hours will not be counted as a new event.
Cardiovascular risk factor modification	
Smoking	Self-reported number of cigarettes
Alcohol intake	Self-reported grams of alcohol consumed/week
Body mass index	Self-reported weight and height
Nutrition	CSIRO Healthy Diet Score ⁴¹
Diabetes control	Hb A1C
Exercise, physical activity and functional capacity	6-minute walk test ⁴² International Physical Activity Questionnaire-Short Form ⁴³

	Specific Activity Questionnaire ⁴⁴
Medication adherence	Adherence in Chronic Diseases Scale ⁴⁵
<i>Patient-reported outcome measures (PROMs) and patient-reported experience measures (PREMs)</i>	
Anxiety	Cardiac anxiety questionnaire ⁴⁶
Depression	PHQ-2, PHQ-9 ⁴⁷
Quality of life*	EQ-5D-5L ⁴⁸ Kansas City Cardiomyopathy Questionnaire-12 ⁴⁹ (for patients with heart failure)
Health literacy	Brief Health Literacy Screening tool ⁵⁰
Patient satisfaction	Self-reported satisfaction

PHQ-2 and PHQ-9: the 2- and 9-question Patient Health Questionnaires, respectively; EQ-5D-5L: European Quality of Life- 5 Dimensions – 5 Levels;

Patients and public involvement

The CHAP model of care is underpinned by a patient-centred approach. Patient-centred care or person-centred care is based on principles that those receiving healthcare must be treated with dignity and respect and be involved in all decisions about their health.⁵¹ This approach is applied in this study through involvement of patients in the co-design of the CR web program⁴⁰ and development of a model that allows patients to choose among different CR modes of delivery (face-to-face, telephone or web-based, or combinations of these modes- Figure 2).^{19 51}

⁵² We will particularly promote the participation of older CR patients and people from cultural and linguistically diverse backgrounds to ensure that groups at increased risk of low technology and health literacy are involved in the co-designing of the web-based CR program. We anticipate this will increase patient adherence and engagement with the web-based CR program.⁵²

Updating the modalities of telehealth that are currently available will allow for greater flexibility of CR delivery which can improve access for patients in need. Importantly, this also

provides an appropriate response to the COVID-19 pandemic when traditional CR service delivery is disrupted.^{15 53 54}

3) Lack of sustainable lifelong commitment to CR supported by primary care

CHAP project will build upon the GP Hybrid Model of CR delivery by creating a business model which allows primary care to support CR delivery and assessment of CR results (risk factor change) of patients undertaking telephone and web-based CR. The model will comprise a 'Heart Health Risk Factor' pre-assessment at the beginning of the program (i.e. 1-2 weeks post-discharge) which will last 75 minutes (60 minutes with a Practice Nurse and 15 minutes with the GP). Re-assessments will occur at 8-12 weeks after the beginning of the program, and 6 months and 12 months post-completion of the CR program. These will comprise 45 to 60 minutes consultations with the practice nurse, and 15-minute consultations with the GP. This has potential to enable long term support for heart health from GPs.

4) Heterogeneous CR quality

The CHAP project will seek adherence to the quality indicators through multiple online and offline professional development programs targeting the practitioners, development of an accreditation program and refinement of data collection tools on the national CR quality indicators. In particular, the project will try to integrate objective patient-generated physical activity tracking data to the CATCH CR database to enable objective monitoring of physical activity during CR and personalised feedback.

Outcomes

Primary outcome – CR attendance rate defined as the number of patients commencing CR (\geq 1 session) among those eligible to CR.

Secondary outcomes - will compare risk factor modification, clinical outcomes, PROMS and PREMS of the CHAP and non-CHAP groups. Measurements, tools and questionnaires and data

sources for the secondary outcomes are presented in Table 1. The selection of validated tools and questionnaires followed extensive consultations with CR clinicians about their preferences and implications of the tools to documentation time and costs to the services.

Data Sources

The South Australian CR clinical database (the CATCH database) collects data uniformly across metropolitan, rural and remote areas using a standardised electronic entry process, including electronic data capture, of patients moving through public CR services from rural and remote to a metropolitan hospital or vice versa.¹⁹ This database will be linked to the SA Health administrative database, enabling follow-up and assessment of hospital re-admissions, Emergency Department (ED) presentations and mortality.⁵⁵ This data linkage approach is similar to the one used to audit the CR services in South Australia from 2013-2015.¹²

Bias

To avoid information bias (i.e., systematic differences in the completeness or the accuracy of data leading to differential misclassification of individuals regarding exposures or outcomes) we are up-skilling clinicians to input data onto the CR databases as part of continuous quality improvement processes. The goal is to fill the gap of scarcity of data on CR data in Australia⁵⁶ and achieve universal, uniform data entry in South Australian CR programs.

Sample size

The average state-wide attendance of CR from 2013-2015 was 28.3% (pre-CHAP estimate)¹⁰. Based upon a 4% improvement in attendance (i.e., to 32.3%) as reported by the British Association for Cardiovascular Prevention and Rehabilitation audit,⁵⁷ a total sample size of 1223 (CHAP) and 3669 (usual care, 1:3 matching) will produce a two-sided 95% confidence interval for the difference in population proportions with a width that is equal to 0.06. Allowing

for a 20% dropout, we will need to observe 1529 patients (765 per year) in the group exposed to CHAP that will be matched to 4587 patients in the group non-exposed to CHAP (usual care) from the CATCH database. From 2015-2019, approximately 2000 patients were entered into the CATCH CR database per year making this study highly feasible.

Data analysis

To evaluate the effectiveness of CHAP (Objective 2)

Unpaired *t*-tests, χ tests and Mann–Whitney *U* tests for non-normal data will be used for comparisons for the pre-post analysis of the CHAP model components. Statistical significance will be inferred if 2-tailed tests estimated $P < 0.05$. Rates of referral, attendance, and completion of CR per year; modification of risk factors; adherence to evidence-based pharmacotherapy morbidity and mortality at 30 days and 12 months will be compared between CHAP and non-CHAP attendees and presented as Odds Ratios and 95% Confidence Intervals. Matching will be conducted by grouping all cases and controls into relevant groupings of Age, Sex and DRG and analysed using the Mantel-Haenszel methods.⁵⁸ Cox regression models adjusted for clinical and social factors (age, sex, the Index of Relative Socio-Economic Advantage and Disadvantage (IRSAD) from the Australian Bureau of Statistics' Socio-Economic Indexes for Areas-SEIFA score,⁵⁹ accessibility and Remoteness Index of Australia (ARIA+),⁶⁰ primary cardiac diagnosis, Charlson Index,⁶¹ in-hospital length of stay, prior heart failure, coronary disease, atrial fibrillation, revascularisation, malignancy) will be performed to understand the association between exposure to the CHAP model and attendance. For clinical outcomes (30-day and 12-month hospital CVD-related hospital re-admission, 30-day and 12-month CVD-related mortality), the model will also be adjusted for CR attendance.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

To evaluate the cost-effectiveness of CHAP (Objective 3)

The economic evaluation of the CHAP model including the business model (explore the item numbers that can be utilised to cover GP based CR and items that can be used to employ and train practice nurses to assist with assessment, monitoring, collection of outcomes data and triage to maximise GP consultation time) that involves primary care in CR delivery will comprise: i) the cost efficiencies to health services, and ii) cost-benefit to rural and remote patients and families.

i. The cost efficiencies to the health services

All CVD hospitalisations during the follow-up period will be included and assigned a cost, based on the diagnosis-related group category reference and length of stay. Costs of providing CR will be taken from a health system perspective and calculated over 1 year; this includes staffing, consumables, overheads, and capital expenses. Quality-of-life measures will be estimated at baseline, 6 and 12 months after CR completion, using the standard EQ-5D-5L.⁴⁸ The utility is defined by the Department of Health and Ageing as a numerical value assigned by an individual to a preference for, or desirability of, a specific level of health status or a specific health outcome. Data on utilities and survival can be combined to calculate Quality-Adjusted Life-Year (QALY) by exposed and non-exposed groups using the Quality-Adjusted Survival Analysis (QASA) method.⁶²⁻⁶⁴ Incremental cost per QALY saved will be calculated as an incremental (rehabilitation v usual care) cost per patient divided by the incremental QALY per patient.⁶⁵

ii. Cost-benefit to rural and remote patients and families

Using an economic survey developed by the Western Australian Department of Health⁶⁶ we will quantify the economic impact of the CHAP model based on information at baseline including patient demographics such as household structure, employment status and the

distances travelled to access services; the nature of the consultation, waiting time, whether the patient required time off work (and if so, how much), and the mode of transport used to travel to the hospital. The survey will include accompanying persons, and if the person required any time off work to do so. Finally, the survey will explore the patient's treatment options; for example, did they consider having an appointment in the city essential to their care? This data will be modelled to compare two scenarios for face-to-face CR; 1) if the patient travelled to the city to attend CR, or 2) if the patient travelled to the nearest rural CR program if this is within 60 minutes' drive (approximately 100 km). All results will be presented unadjusted for multiple comparisons and all analyses will be done on an intention-to-treat basis. 95% confidence intervals for the difference in QALYs will be calculated with bootstrapping.⁶³

Time Frame of the Project

The original timeframe comprised a three-year project from 1st July, 2019 to 30th June, 2022. Due to the impact of the COVID pandemic on the project implementation, the project had a 12-month extension and will now be complete by 30th June 2023. Year 1-2 will include project staff employment and training, development of resources, education materials, study protocols, additional datasets, and Ethics approval. Recruitment will be conducted over years 2-3 and follow-up over years 3-4.

ETHICS AND DISSEMINATION

Ethics approval and consent to participate

This study was approved by the Southern Adelaide Clinical Human Research Ethics Committee Research Committee on 28th, April 2020 (HREC/20/SAC/78). The data linkage between the South Australian CR clinical database and the Department of Health

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

administrative databases was approved by the Department for Health and Wellbeing on 26th August, 2021 (2021/HRE00270). All consecutive patients who are eligible for CR according to the recommended DRG codes and referred for CR will have consent waiver to have their data used for this research.

Dissemination

Findings will be disseminated to patients and health professionals using various means and media. A public website (<https://chaproject.com.au>), ³⁶ online educational sessions with patients and health professionals working in CR, monthly newsletters sent to participants via email, social media (Twitter @CHAPproject, LinkedIn CHAP Project and Facebook), and scientific publications and conference presentations.

De-identified data will be available upon reasonable request.

Authors contributions: The study was conceived and designed by RC, SN, AB, DC, JB, AM, CM, VV, JH, PT, BK, CZ, IP, RT, RT, and WK. The first draft of the paper was written by AB with significant input from RC before receiving input from remaining authors; CA, KG, KN, HD, SC, MAPP, IL, VP, MR, KW, HM, AK, LG, ML and CG. All authors read and approved the final manuscript.

Funding: This study is supported by a peer-reviewed National Health and Medical Research Council (NHMRC Partnership Grant-GNT1169893) and by the following partners: Heart Foundation, Exercise and Sports Science Australia (ESSA), Australian Cardiac Rehabilitation Association (ACRA), Country SA Primary Health Network, Cardiac Society of Australia and New Zealand (CSANZ), Flinders Foundation, and Royal College of General Practitioners (RACGP), Astra Zeneca, Novartis, Health Translation SA, the Integrated Cardiac Clinical Network (iCCnet SA) and Country Health SA and the Enterprise Data and Information Department of Health and Wellbeing, SA Health.

Competing interests: The authors do not have competing interests.

Acknowledgements: VLV is funded by the Australian Government Department of Health through the Rural Health Multidisciplinary Training (RHMT) program.

JMH is supported by a Future Leader Fellowship provided by the National Heart Foundation of Australia.

CM is funded by a Medical Research Future Fund Investigator Grant (1193862).

1
2
3 431 **Legend of Figures and Tables**
4

5 432
6
7 433 **Figures**
8

9 434 Figure 1: Using the Model for Large Scale Knowledge Translation to develop CHAP

10
11 435 Figure 2: Addressing the barriers through the CHAP model of care (Exposure) according to the
12 436 4th Stage of the Model for Large Scale Knowledge Translation (‘Engage, Educate, Execute,
13 437 Evaluate’)
14 438

15
16 439

17
18
19 440

20
21
22 441

23
24 442
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Cardiovascular diseases - WHO.: World Health Organization
2020 [Available from: https://www.who.int/health-topics/cardiovascular-diseases/#tab=tab_1
accessed 19/05/2021 2021.
2. Causes of Death 2019. ABS cat no 33030: Australian Bureau of Statistics (ABS), 2020.
3. Heart, stroke & vascular diseases: Australian Institute of Health & Welfare (AIHW), 2018.
4. Clark RA, Coffee N, Turner D, et al. Application of geographic modeling techniques to quantify spatial
access to health services before and after an acute cardiac event: the Cardiac Accessibility and
Remoteness Index for Australia (ARIA) project. *Circulation* 2012;125(16):2006-14. doi:
10.1161/circulationaha.111.083394 [published Online First: 2012/03/28]
5. Australia NHFo. Interactive Australian Heart Maps. Compare heart health indicators across
Australia. 2021 [Available from: [https://www.heartfoundation.org.au/health-professional-
tools/interactive-heart-map-australia](https://www.heartfoundation.org.au/health-professional-tools/interactive-heart-map-australia) accessed 16/06/2021 accessed 16/06/2021 2021.
6. Tideman P, Taylor AW, Janus E, et al. A comparison of Australian rural and metropolitan
cardiovascular risk and mortality: the Greater Green Triangle and North West Adelaide
population surveys. *BMJ Open* 2013;3(8):e003203. doi: 10.1136/bmjopen-2013-003203
7. Statistics ABo. 2016 Census Quick Stats. South Australia 2016 [Available from:
[https://quickstats.censusdata.abs.gov.au/census_services/getproduct/census/2016/quicksta
t/4?opendocument](https://quickstats.censusdata.abs.gov.au/census_services/getproduct/census/2016/quickstat/4?opendocument) accessed 16/06/2021 2021.
8. A time for action in South Australia. : National Heart Foundation of Australia
2018.
9. Woodruffe S, Neubeck L, Clark RA, et al. Australian Cardiovascular Health and Rehabilitation
Association (ACRA) core components of cardiovascular disease secondary prevention and
cardiac rehabilitation 2014. *Heart Lung Circ* 2015;24(5):430-41. doi:
10.1016/j.hlc.2014.12.008 [published Online First: 2015/02/01]
10. Anderson L, Thompson DR, Oldridge N, et al. Exercise-based cardiac rehabilitation for coronary
heart disease. *Cochrane Database Syst Rev* 2016 Jan 5;2016(1):CD001800 doi:
101002/14651858CD001800pub3 PMID: 26730878; PMCID: PMC6491180 2016
11. Redfern J, Hyun K, Chew DP, et al. Prescription of secondary prevention medications, lifestyle
advice, and referral to rehabilitation among acute coronary syndrome inpatients: results from
a large prospective audit in Australia and New Zealand. *Heart* 2014;100(16):1281-8. doi:
10.1136/heartjnl-2013-305296 [published Online First: 2014/06/09]
12. Astley CM, Chew DP, Keech W, et al. The Impact of Cardiac Rehabilitation and Secondary
Prevention Programs on 12-Month Clinical Outcomes: A Linked Data Analysis. *Heart Lung Circ*
2020;29(3):475-82. doi: 10.1016/j.hlc.2019.03.015 [published Online First: 2019/05/11]
13. Jelinek MV, Thompson DR, Ski C, et al. 40 years of cardiac rehabilitation and secondary prevention
in post-cardiac ischaemic patients. Are we still in the wilderness? *Int J Cardiol* 2015;179:153-
9. doi: 10.1016/j.ijcard.2014.10.154 [published Online First: 2014/12/03]
14. Clark RA, Conway A, Poulsen V, et al. Alternative models of cardiac rehabilitation: a systematic
review. *Eur J Prev Cardiol* 2015;22(1):35-74. doi: 10.1177/2047487313501093 [published
Online First: 2013/08/15]
15. Dalal HM, Doherty P, McDonagh STJ, et al. Virtual and in-person cardiac rehabilitation. *BMJ*
2021;373:n1270. doi: 10.1136/bmj.n1270
16. West RR, Jones DA, Henderson AH. Rehabilitation after myocardial infarction trial (RAMIT): multi-
centre randomised controlled trial of comprehensive cardiac rehabilitation in patients
following acute myocardial infarction. *Heart* 2012;98(8):637-44. doi: 10.1136/heartjnl-2011-
300302 [published Online First: 2011/12/24]

17. Astley CM, Neubeck L, Gallagher R, et al. Cardiac Rehabilitation: Unraveling the Complexity of Referral and Current Models of Delivery. *J Cardiovasc Nurs* 2017;32(3):236-43. doi: 10.1097/jcn.0000000000000332 [published Online First: 2016/04/15]
18. Tideman PA, Tirimacco R, Senior DP, et al. Impact of a regionalised clinical cardiac support network on mortality among rural patients with myocardial infarction. *Med J Aust* 2014;200(3):157-60. doi: 10.5694/mja13.10645 [published Online First: 2014/02/18]
19. Tideman P, Tirimacco R, Berry N, et al. Country access to cardiac health (CATCH) program: improving rehospitalisation rates and length of stay for cardiac rehabilitation using innovative telehealth delivery. *Heart, Lung and Circulation* 2015;24:S453. doi: 10.1016/j.hlc.2015.06.790
20. Clark RA, Wilkinson D, Coffee N, et al. Mapping services to support a patient's journey through evidence-based care pathways after a cardiac event. *Heart, Lung and Circulation* 2009;18:S266. doi: 10.1016/j.hlc.2009.05.656
21. Gallagher R, Neubeck L, Du H, et al. Facilitating or getting in the way? The effect of clinicians' knowledge, values and beliefs on referral and participation. *Eur J Prev Cardiol* 2016;23(11):1141-50. doi: 10.1177/2047487316630085 [published Online First: 2016/02/03]
22. Ghisi GL, Polyzotis P, Oh P, et al. Physician factors affecting cardiac rehabilitation referral and patient enrollment: a systematic review. *Clin Cardiol* 2013;36(6):323-35. doi: 10.1002/clc.22126 [published Online First: 2013/05/04]
23. Summit to address rural medical specialist shortage: Royal Australian College of General Practitioners 2018 [Available from: <https://www1.racgp.org.au/newsgp/professional/summit-to-address-rural-medical-specialist-shortage> accessed 18/01/2021 2021.
24. Clark C, Tirimacco R, Morfidis N, et al. Delivery of outpatient cardiac rehabilitation using a GP Hybrid/Telephone Program model. National Rural Health Alliance. Tasmania: Rural Health Org 2019.
25. Pack QR, Johnson LL, Barr LM, et al. Improving cardiac rehabilitation attendance and completion through quality improvement activities and a motivational program. *J Cardiopulm Rehabil Prev* 2013;33(3):153-9. doi: 10.1097/HCR.0b013e31828db386 [published Online First: 2013/04/19]
26. Gallagher R, Thomas E, Astley C, et al. Cardiac Rehabilitation Quality in Australia: Proposed National Indicator for Field-Testing. *Heart, Lung and Circulation* 2020;29(9):1273-77. doi: 10.1016/j.hlc.2020.02.014
27. Peters DH, Adam T, Alonge O, et al. Implementation research: what it is and how to do it. *BMJ : British Medical Journal* 2013;347:f6753. doi: 10.1136/bmj.f6753
28. Pronovost PJ, Berenholtz SM, Needham DM. Translating evidence into practice: a model for large scale knowledge translation. *BMJ* 2008;337:a1714. doi: 10.1136/bmj.a1714
29. Abreu A, Frederix I, Dendale P, et al. Standardization and quality improvement of secondary prevention through cardiovascular rehabilitation programmes in Europe: The avenue towards EAPC accreditation programme: A position statement of the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology (EAPC). *European Journal of Preventive Cardiology* 2020;2047487320924912. doi: 10.1177/2047487320924912
30. Pearce N. Analysis of matched case-control studies. *BMJ* 2016;352:i969. doi: 10.1136/bmj.i969
31. Country SA Primary Health Network SA Health 2021 [Available from: <https://www.countrysaphn.com.au/> accessed 19/05/2021 2021.
32. Chew DP, Scott IA, Cullen L, et al. National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand: Australian clinical guidelines for the management of acute coronary syndromes 2016. *Med J Aust* 2016;205(3):128-33. doi: 10.5694/mja16.00368 [published Online First: 2016/07/29]
33. International Statistical Classification of Diseases and Related Health Problems 10th Revision: World Health Organization, 2016.
34. Independent Hospital Pricing Authority. International Classification of Diseases and Related Health Problems, 11th revision. Australian Modification (ICD-10-AM), 2019.

35. ABS. Census of Population and Housing: Reflecting Australia - Series form the Census, 2016. Aboriginal and Torres Strait Islander Population, 2016. 2017 [Available from: <https://www.abs.gov.au/statistics/people/aboriginal-and-torres-strait-islander-peoples/estimates-aboriginal-and-torres-strait-islander-australians/latest-release> accessed 16/12/2021 2021.
36. Country Heart Attack Prevention Project 2021 [Available from: chapproject.com.au accessed 16/12/2021 2021.
37. Norman CD, Skinner HA. eHealth Literacy: Essential Skills for Consumer Health in a Networked World. *J Med Internet Res* 2006;8(2):e9. doi: 10.2196/jmir.8.2.e9
38. Integrated Point of Care Clinical System Integrated Clinical Cardiovascular Network (iCCnet) SA 2021 [Available from: <https://www.iccnetsa.org.au/ipoccs.aspx> accessed 19/05/2021 2021.
39. Will T. User Experience (UX): Process and Methodology 2020 [Available from: <https://uiuxtrend.com/user-experience-ux-process/>.
40. Nesbitt K, Beleigoli A, Du H, et al. Web-Based Cardiac Rehabilitation: A Co-Design Workshop. *Telehealth Innovations in Remote Healthcare Services Delivery* 2021; 277. <https://ebooks.iospress.nl/volumearticle/56656>.
41. Hendrie GA, Rebuli MA, Golley RK. Reliability and relative validity of a diet index score for adults derived from a self-reported short food survey. *Nutr Diet* 2017;74(3):291-97. doi: 10.1111/1747-0080.12303 [published Online First: 2017/07/22]
42. Bellet RN, Adams L, Morris NR. The 6-minute walk test in outpatient cardiac rehabilitation: validity, reliability and responsiveness—a systematic review. *Physiotherapy* 2012;98(4):277-86. doi: <https://doi.org/10.1016/j.physio.2011.11.003>
43. Craig CL, Marshall AL, Sjöström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95. doi: 10.1249/01.Mss.0000078924.61453.Fb [published Online First: 2003/08/06]
44. Rankin SL, Briffa TG, Morton AR, et al. A specific activity questionnaire to measure the functional capacity of cardiac patients. *Am J Cardiol* 1996;77(14):1220-3. doi: 10.1016/s0002-9149(97)89157-6 [published Online First: 1996/06/01]
45. Buszko K, Obońska K, Michalski P, et al. The Adherence Scale in Chronic Diseases (ASCD). The power of knowledge: the key to successful patient — health care provider cooperation. *Medical Research Journal* 2016;1(1):37-42. doi: 10.5603/MRJ.2016.0006
46. Eifert GH, Thompson RN, Zvolensky MJ, et al. The cardiac anxiety questionnaire: development and preliminary validity. *Behav Res Ther* 2000;38(10):1039-53. doi: 10.1016/s0005-7967(99)00132-1 [published Online First: 2000/09/27]
47. Arroll B, Goodyear-Smith F, Crengle S, et al. Validation of PHQ-2 and PHQ-9 to screen for major depression in the primary care population. *Ann Fam Med* 2010;8(4):348-53. doi: 10.1370/afm.1139 [published Online First: 2010/07/21]
48. McCaffrey N, Kaambwa B, Currow DC, et al. Health-related quality of life measured using the EQ-5D-5L: South Australian population norms. *Health Qual Life Outcomes* 2016;14(1):133. doi: 10.1186/s12955-016-0537-0 [published Online First: 2016/09/21]
49. Spertus JA, Jones PG. Development and Validation of a Short Version of the Kansas City Cardiomyopathy Questionnaire. *Circulation: Cardiovascular Quality and Outcomes* 2015;8(5):469-76. doi: 10.1161/CIRCOUTCOMES.115.001958
50. Haun J, Luther S, Dodd V, et al. Measurement variation across health literacy assessments: implications for assessment selection in research and practice. *J Health Commun* 2012;17 Suppl 3:141-59. doi: 10.1080/10810730.2012.712615 [published Online First: 2012/10/04]
51. Patient-Centred Care Improving Quality and Safety Through Partnerships With Patients and Consumers. *Australia Commission on Safety and Quality in Healthcare* 2011
52. Safety and Quality Improvement Guide Standard 2: Partnering with Consumers. In: Australia Co, ed. Sydney: Australia Commission on Safety and Quality in Healthcare, 2012.

1
2
3 592 53. Nicholls SJ, Nelson M, Astley C, et al. Optimising Secondary Prevention and Cardiac Rehabilitation
4 593 for Atherosclerotic Cardiovascular Disease During the COVID-19 Pandemic: A Position
5 594 Statement From the Cardiac Society of Australia and New Zealand (CSANZ)☆. *Heart, Lung and*
6 595 *Circulation* 2020;29(7):e99-e104. doi: <https://doi.org/10.1016/j.hlc.2020.04.007>
7 596 54. Arnold RH, Tideman PA, Devlin GP, et al. Rural and Remote Cardiology During the COVID-19
8 597 Pandemic: Cardiac Society of Australia and New Zealand (CSANZ) Consensus
9 598 Statement^{☆}. *Heart, Lung and Circulation* 2020;29(7):e88-e93. doi:
10 599 10.1016/j.hlc.2020.05.001
11 600 55. Admitted Patient Care Government of South Australia SA Health 2021 [accessed 18/01/2021 2021.
12 601 56. Driscoll A, Hinde S, Harrison A, et al. Estimating the health loss due to poor engagement with
13 602 cardiac rehabilitation in Australia. *International Journal of Cardiology* 2020;317:7-12. doi:
14 603 <https://doi.org/10.1016/j.ijcard.2020.04.088>
15 604 57. National Audit of Cardiac Rehabilitation (NACR) Quality and Outcomes Report 2018: British Heart
16 605 Foundation, 2018.
17 606 58. Altman D. Practical Statistics for Medical Research. London: Chapman and Hall 1991.
18 607 59. Census of Population and Housing: Socio-economic Indexes for Areas (SEIFA): Australian Bureau
19 608 of Statistics; 2016 [Available from:
20 609 <https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/2033.0.55.001main+features12016>
21 610 accessed 18/01/2021 2021.
22 611 60. Australian Statistical Geography Standard: Volume 5 - Remoteness Structure.: Australian Bureau
23 612 of Statistics; 2018 [Available from:
24 613 <https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.005> accessed 18/01/2021 2021.
25 614 61. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in
26 615 longitudinal studies: development and validation. *J Chronic Dis* 1987;40(5):373-83. doi:
27 616 10.1016/0021-9681(87)90171-8 [published Online First: 1987/01/01]
28 617 62. Glasziou PP, Simes RJ, Gelber RD. Quality adjusted survival analysis. *Stat Med* 1990;9(11):1259-76.
29 618 doi: 10.1002/sim.4780091106 [published Online First: 1990/11/01]
30 619 63. Glasziou PP, Cole BF, Gelber RD, et al. Quality adjusted survival analysis with repeated quality of
31 620 life measures. *Stat Med* 1998;17(11):1215-29. doi: 10.1002/(sici)1097-
32 621 0258(19980615)17:11<1215::aid-sim844>3.0.co;2-y [published Online First: 1998/07/22]
33 622 64. Martin AJ, Glasziou PP, Simes RJ. A cardiovascular extension of the Health Measurement
34 623 Questionnaire. *J Epidemiol Community Health* 1999;53(9):548-57. doi: 10.1136/jech.53.9.548
35 624 [published Online First: 1999/11/24]
36 625 65. Sullivan SD, Mauskopf JA, Augustovski F, et al. Budget impact analysis-principles of good practice:
37 626 report of the ISPOR 2012 Budget Impact Analysis Good Practice II Task Force. *Value Health*
38 627 2014;17(1):5-14. doi: 10.1016/j.jval.2013.08.2291 [published Online First: 2014/01/21]
39 628 66. Rankin S, House A, Heath D. The real cost of accessing specialist medical services for rural and
40 629 remote communities in Western Australia. . 5th National Rural Health Conference 1999.
41 630
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Using the Model for Large Scale Knowledge Translation to develop CHAP

276x126mm (150 x 150 DPI)




Figure 2: Addressing the barriers through the CHAP model of care (Exposure) according to the 4th Stage of the Model for Large Scale Knowledge Translation ('Engage, Educate, Execute, Evaluate')

174x191mm (96 x 96 DPI)

STROBE CHECKLIST

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	1 4
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	8-10
Objectives	3	State specific objectives, including any prespecified hypotheses	11
Methods			
Study design	4	Present key elements of study design early in the paper	12
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	13
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	14
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	14-18
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	18
Bias	9	Describe any efforts to address potential sources of bias	19
Study size	10	Explain how the study size was arrived at	19

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	20-21 Table 1
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	20-21
Results -Not applicable			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
<hr/>		
Discussion	Not applicable	
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
<hr/>		
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based Page 23

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only